



UNIVERSITÀ DEGLI STUDI DI SASSARI
FACOLTÀ DI MEDICINA E CHIRURGIA
Scuola di Dottorato in Scienze Biomediche
Direttore Prof. Eusebio Tolu

Indirizzo in Odontostomatologia Preventiva
Direttore Prof.ssa Egle Milia

EARLY CHILDHOOD CARIES (ECC) IS STILL A PROBLEM? AN
EPIDEMIOLOGICAL AND AETIOLOGICAL APPROACH.

Tutor:
Prof.ssa PierFranca Lugliè

Tesi di:
Dr.ssa Giovanna Congiu

Introduzione

La prevalenza della patologia cariosa nei paesi occidentali è diminuita in maniera decisa negli ultimi 20 anni, nonostante ciò, all'interno di tali società, esistono gruppi a rischio quali i bambini e gli anziani (Curzon e Preston, 2004). Per anni la patologia cariosa nell'età infantile è stata descritta con vari termini quali: carie da biberon, carie rampante o carie infantile (De Grauwe e Martens, 2004). Attualmente, si preferisce utilizzare il termine Early Childhood Caries (ECC), coniato dall'American Academy of Paediatric Dentistry (AAPD, 2008), per descrivere la presenza di almeno un elemento dentario cariato, otturato o estratto per carie nella dentatura decidua. Tale termine comunemente accettato nella letteratura più recente come definizione più generale include tra le cause della patologia sia l'allattamento al seno sia l'uso di sostanze edulcorate (De Grauwe e Martens, 2004).

Il quadro clinico dell'ECC è caratterizzato da lesioni cariose multifocali e gravi a livello delle superfici vestibolari e palatali degli incisivi superiori e di quelle palatali dei molari superiori. Il processo carioso ha inizio 3-6 mesi dopo l'eruzione dei denti interessando gli incisivi superiori e raggiunge l'apice prima dei 18 mesi (Thitasomakul *et al.*, 2006). Gli incisivi inferiori sono coinvolti solo tardivamente dal processo carioso e nei casi estremamente gravi grazie alla loro posizione vicino ai dotti escretori delle ghiandole sottomandibolari e sottolinguali che determina una continua detersione delle superfici. Inoltre, la posizione d'appoggio della lingua su tali elementi dentari durante la suzione ha una funzione protettiva permettendo a una minore quantità di cibo di aderire alle superfici dentali (De Grauwe e Martens, 2004).

Abitudini alimentari scorrette e l'allattamento al seno prolungato, sia in termini di ore giornaliere sia complessivamente oltre i 12 mesi, sono direttamente correlati all'insorgenza dell'ECC (Campus *et al.*, 2007; Al-Dashti *et al.*, 1995; Olak *et al.*, 2007).

Non si deve, infine, sottovalutare il potenziale cariogeno dei farmaci in uso in pediatria; ai piccoli pazienti sono somministrati regolarmente, in corso di malattie sistemiche, sciroppi zuccherati o

medicinali che riducono la salivazione (nei pazienti asmatici) (Horowitz, 1998; Curzon e Preston, 2004).

Fattore rilevante nell'eziologia dell'ECC è lo stato socio-economico. Da vari studi effettuati nelle scuole dell'obbligo è risultato che la distribuzione della carie è concentrata maggiormente nei quartieri popolari dove vivono immigrati e famiglie con problemi socio-economici. Dai questionari compilati dai genitori è emerso che la maggior parte delle famiglie è composta da tre o più figli, vive con lo stipendio del padre, e sia il padre che la madre, spesso di giovane età, hanno conseguito un livello primario d'istruzione e in gran parte sono extracomunitari. Dal punto di vista odontoiatrico gran parte dei genitori hanno dichiarato di non conoscere le misure di prevenzione primaria della carie, di aver avuto esperienze di patologia cariosa e di ricorrere alle cure odontoiatriche solo in caso di urgenze. L'inconsapevolezza di avere problemi dentali, l'assenza di mal di denti, la spesa troppo alta per le cure odontoiatriche, la paura del dentista e la perdita dei rimborsi per le cure effettuate sono le ragioni più citate in merito alla scarsità delle visite odontoiatriche (Martens *et al.*, 2006; Azogui-Lèvy *et al.*, 2003; Donaldson et Kinirons, 2001; Blair *et al.*, 2006).

Con tale tesi si è voluto analizzare il problema dell'ECC sotto diversi aspetti: eziologico, epidemiologico e preventivo.

Il lavoro è stato diviso in tre fasi:

1. revisione sistematica della letteratura sulla prevalenza dell'ECC e dei fattori di rischio;
2. studio epidemiologico cross-sectional sulla prevalenza di ECC e fattori di rischio nella popolazione prescolare della città di Sassari.
3. studio clinico randomizzato sull'efficacia della somministrazione nelle donne, durante l'allattamento, di un integratore contenente fluoro.

References

- Al-Dashti AA, Williams SA, Curzon ME. Breast feeding, bottle feeding and dental caries in Kuwait, a country with low-fluoride levels in water supply; *Community Dental Health* 1995;12:42-47.
- American Academy of Paediatric Dentistry (2008). Definition of Early Childhood Caries (ECC). http://www.aapd.org/media/policies_guidelines/d_ecc.pdf
- Azogui-Lèvy S, Lombrail P, Riordan PJ, Brodin M, Baillon-Javon E, Pirlet MC, Boy-Lefèvre ML. Evaluation of dental care program for school beginners in a Paris suburb; *Community Dental Oral Epidemiol* 2003;31:285-291.
- Blair Y, Macpherson L, McCall D, McMahon A. Dental health of 5 years old following community-based oral health promotion in Glasgow; *International Journal of Paediatric Dentistry* 2006;16:388-398.
- Campus G, Solinas G, Sanna A, Maida C, Castiglia P. Determinants of ECC in Sardinian preschool children. *Community Dent Health* 2007;24:253-6.
- Curzon ME, Preston AJ. Risk groups: nursing bottle caries/caries in the elderly. *Caries Res* 2004;38:24-33.
- De Grauwe A, Aps JK, Martens LC. Early Childhood Caries (ECC): what's in a name? *Eur J Paediatr Dent* 2004;5:62-70.
- Donaldson M, Kinirons M. The effectiveness of the school dental screening programme in stimulating dental attendance for children in need of treatment in Northern Ireland; *Community Dent Oral Epidemiol* 2001; 29:143-149.
- Horowitz HS. Research issues in Early Childhood Caries. *Community Dent Oral Epidemiol* 1998;26:67-81.

Martens L, Vannobbergen J, Willems S, Aps J, De Maeseneer J. Determinants of early childhood caries in a group of inner-city children; Quintessence International 2006;37:527-536.

Olak J, Mändar R, Karjalainen S, Söderling E, Saag M. Dental health and oral mutans streptococci in 2-4 years old Estonian children; International Journal of Paediatric Dentistry 2007;17:92-97.

Thitasomakul S, Thearmontree A, Piwat S, Chankanka O, Pithpornchaiyakul W, Teanpaisan R, Madyusoh S.; A longitudinal study of early childhood caries in 9- to 18-month-old Thai infants. Community Dent Oral Epidemiol 2006;34:429-436.

PAPER I

Early Childhood Caries (ECC) prevalence and background factors. A review.

Congiu Giovanna^a, Lugliè PierFranca^a, Campus Guglielmo^a.

^aDental Institute, University of Sassari.

Corresponding author
Dr Guglielmo Campus
Dental Institute, University of Sassari
Viale san Pietro 43/C I-07100 Sassari, Italy
Tel.+39-079-228540
Fax. +39-079-228541
Email gcampus@uniss.it

Summary

Aim: to describe the prevalence of ECC by reviewing published reports, and secondly to assess the rule of background factors on the ECC prevalence.

Design: Included studies were: reviews, meta analysis, RCT assessing the prevalence of ECC including more than 100 subjects; reporting subjects without any stated medical condition; in vivo studies; issued from January 1st, 2000 to May 31st, 2011. The age range for the inclusion was 0 to five years of age.

Results: Database searching showed 411 findings as potentially eligible. After abstract review 8 papers were retrieved as full-text and assessed for eligibility: three using ECC as mesh word, four using the term Early Childhood caries, and one using S-ECC as mesh word.

Conclusions: The main risk factor for ECC is the low socio-economical level of the parents. ECC is a public health problem and it requires involvement of all health professionals that provide care to children together with efforts from family members.

The dental caries is still an endemic disease affecting especially young subjects. Dental caries is a preventable disease, and it can be stopped and even potentially reversed during its early stages. People remain susceptible to the disease throughout their lives. Children in the age range of 12–30 months have a special caries pattern that differs from that in older children. The disease develops and affects the maxillary primary incisors and first primary molars in a way that reflects the pattern of eruption, longer the tooth has been present and exposed to the caries challenge, the more it is affected. So the teeth in the upper jaw are most vulnerable, while teeth in mandibular are protected by the tongue and by saliva from submandibular and sublingual glands. Caries lesion in young child is a worsening form of decay. Whereas certain feeding patterns, such as “at will” breast feeding and bedtime bottle feeding, have long been known to be risk factors for ECC, current research has demonstrated that other factors may also play a role.

This pattern of dental caries has been labeled variously as “bottle caries”, “nursing caries”, “baby bottle tooth decay”, or “night bottle mouth”. These terms suggest that the prime cause of dental caries in early childhood is inappropriate bottle-feeding. Current evidence suggests that use of a sugar-containing liquid in a bottle at night may be an important etiological factor, although it is not necessarily the only etiological factor. Therefore, it is recommended that the term Early Childhood Caries (ECC) be used when describing any form of caries in infants and preschool children. Various systems of classification have been used to define Early Childhood Caries (ECC) (Drury *et al.*, 1999; Psoter *et al.*, 2003). Two commonly accepted classifications for ECC include simple ECC and severe ECC, as defined by the American Academy of Paediatric Dentistry (AAPD, 2008). These two definitions were used for this study as follows:

- Simple ECC any deciduous presence of 1 or more decayed, missing, or filled tooth surfaces in any primary tooth in a child under 6 years of age;

- Severe ECC any sign of smooth-surface caries in children under 3 years of age; 1 or more cavitated, missing, or filled smooth surfaces in primary maxillary anterior teeth in children between the ages of 3 and 5; or decayed, missing, or filled scores of ≥ 4 (ages 3 to < 4), ≥ 5 (ages 4 to < 5), or ≥ 6 (ages 5 to < 6).

The main objective of this paper is aimed to describe the prevalence of ECC by reviewing published reports, and secondly to assess the rule of background factors on the ECC prevalence.

Methods

Protocol for this review was the PRISMA 2009 checklist (available at www.prisma-statement.org)

Eligibility criteria

Included studies were: reviews, meta analysis, RCT assessing the prevalence of ECC including more than 100 subjects; reporting subjects without any stated medical condition; in vivo studies; issued from January 1st, 2000 to May 31st, 2011. The age range for the inclusion was 0 to five years of age.

Only studies published in English language were considered due to the virtual absence of research published in other languages as resulted from preliminary electronic database searches.

Included studies:

- Assessed prevalence and morbidity outcomes for ECC and background variables;
- Measured dental caries as DMFT and/or DMFS Index according to WHO standards;
- Reported dietary, oral hygiene habits and Socio-economic level of the family;
- Clearly described objective, methods and results, with no significant discrepancies;
- Case reports, case series, outbreak investigations and abstracts were excluded.

Possible outcomes for included studies were:

- Increased risk of developing ECC in lower socio-economical families;
- Increased risk of developing ECC associated with dietary habits;

- Increased risk of developing ECC associated with behavioral and/or oral hygiene habits.

Information sources and search

From March 2011 the following electronic databases were searched: Medline, Embase®, The Cochrane Library and Google Scholar®. Two preliminary searches were conducted in June 2011 to obtain an overall idea of findings and to polish searching terms (MeSH words) and limits. The MeSH words was accessed to identify entry terms and compose the final Boolean search: Caries, ECC, S-ECC, Severe Early Childhood Caries, Baby Bottle Caries associated to epidemiology an background factors.

Time coverage was from January 1st, 2000 to May 31st, 2011. No topic related nor relevant finding resulted from both The Cochrane Library and Google Scholar®; these electronic databases were therefore excluded from final Boolean search.

Final search was conducted on September 29th, 2011. Reference lists of included and relevant papers were reviewed. Abstract was collected for all findings.

Study selection and data collection process

One reviewer screened all collected findings and registered title, author and whole reference in two Excel files (one for included and one for excluded findings, according to eligibility criteria) using a screening guide created on eligibility criteria. Kind of source was registered as reason for exclusion. The same reviewer conducted a new and independent screening after 15 days. Duplicates from different electronic databases were excluded. The full text of all studies judged potentially eligible in at least one screening were retrieved. Then, one reviewer screened the full text for inclusion using a screening guide and a second reviewer screened all findings. When disagreement occurred a third reviewer was consulted.

Data items

One reviewer used a standardized and pilot-tested form to abstract data. Then, a second reviewer screened data abstraction. When disagreement occurred, a third independent reviewer was consulted. Abstracted data were about study design, PICO, population, exposure, outcomes, methodological features, results and funding and were defined through STROBE Statement—checklist of items that should be included in reports of observational studies.

The overall quality of evidence for each outcome was rated using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach.

Results

Figure 1 displayed the flow chart of the study. Database searching showed 411 findings as potentially eligible (appendix 1). After abstract review 8 papers were retrieved as full-text and assessed for eligibility: three using ECC as mesh word, four using the term Early Childhood Caries, and one using S-ECC as mesh word. Apart for this single finding, reference list checking highlighted a large number of duplicates with database searching results.

Table 1 describes characteristic of included studies assessing the prevalence of ECC. One study (Bernabè and Hobdell, 2010) was a review paper on the prevalence of ECC in 22 different countries; results reported confirmed that ECC is strictly related to SES level of the family. The authors observed that dental caries prevalence and levels decrease as the Gini per capita increase. The countries were divided in rich countries (according to World Bank criteria) and poor countries. In rich countries, income inequality is the strongest factor associated to Early Childhood Caries. Further epidemiological data must be gathered through surveys in other regions of the country to support oral health programmes. It also needs to raise the awareness on the diagnosis prevention and treatment of ECC among health care workers including paediatricians, physicians, nurses and midwives. Especially facilities for preventive dental work (Fluoride varnish/gel application, fissure sealants) should be enhanced. Two papers were from North America (Tiberia *et al.*, 2007; Nunn *et*

al., 2009) and reported a quite different prevalence rate 3.0% in US and 74.0% in Canada. The two groups were not comparable. Tiberia *et al.* reported about a sample selected from surveys of parents seeking dental care for children in 5 Canadian dental practices so an important bias in selection is present. On the other hand Nunn *et al.* compared data of ECC prevalence among 1- to 3-year-old children sampled from paediatric clinics in Boston area and compare ECC prevalence among these children with similarly aged children who were surveyed as part of the Third National Health and Nutritional Examination Survey (NHANES III). Overall ECC prevalence in Boston children was 3.0 % compared to a double value of prevalence in children derived from NHANES III (6.3%). Two papers were conducted in BRICS (Brazil, Russia, India, China, South-Africa) countries (Tyagi 2008; Feldens *et al.*, 2010). The two papers depicted a prevalence of 19.2% in India, and 63.0% in Brasil. Tyagi described a cross-sectional study on 813 children aged (2-6 years) living in Davangore (India) and no statistical significant difference was detected among gender. No figures were provided how the sample was selected or if it was representative of the area. Feldens *et al.* performed a birth cohort study on southern Brazilian 4 years olds. 500 infants were recruited at birth and the paper was focus only on S-ECC (Severe- Early Childhood Caries) at 4 years of age (final sample 340 subjects). Twenty-six per cent presented simple-ECC and 37.0 % S-ECC.

Two papers were from Japan (Niji *et al.*, 2010) and Greece (Maragakis *et al.*, 2007). Niji *et al.* performed a cross-sectional investigation on the relation between mother age and behavioral and caries prevalence in children. Overall 646 mother-child pairs were examined. ECC prevalence was 41.0%. In Greece, a cross-sectional study on ECC, location and treatment needs was completed. ECC prevalence recorded was quite high (60.1%). A longitudinal observational community-based study was conducted in Thailand (Thitasomakul *et al.*, 2009) using negative binomial regression procedure to analyze data. Overall mean dmfs ranged from 0.1 ± 0.4 to 2.8 ± 2.7 .

Screening for dental caries should start as soon as the first primary tooth erupts or not later than one year of age. Oral health programmes should be established focusing on mothers, caregivers, community health workers, preschool teachers and children.

Table 2 describes the background factors associated with ECC. The paper of Maragakis *et al.* was excluded, because no association with background factors was made. The socio economical level of the family (recorded using several proxy variables like the Gini coefficient, the Annual household income, low income of the family) was reported associated with ECC in four papers (Thitasomakul *et al.*, 2009; Nunn *et al.*, 2009; Barnabé *et al.*, 2010; Feldens *et al.*, 2010). Dietary and behavioral habits like breast-feeding habits, sweetened food, bottle-feeding at night, holding liquids in the mouth for prolonged time, frequency of between meal snacks more than 4 time a day, mother's age at childbirth were positively associated to ECC in six papers (Tiberia *et al.*, 2007; Tyagi, 2008; Nij *et al.*, 2010; Thitasomakul *et al.*, 2009; Feldens *et al.*, 2010).

Discussion

The prevention and management of Early Childhood Caries (ECC) should be an important objective of public health. The onset of the disease is in children with less 2 years of age. The prevention should begin during the prenatal period, continuing with the mother and the child during preschool in collaboration with paediatrician, paediatric dentist and teacher. The vertical transmission (from mother to the child) of cariogenic bacteria involves women in all kind of prevention programmes. This review was aimed to describe and analyze the prevalence of ECC and relative risk factors.

A lot of risk factors are described associated with ECC. However, the main risk factor for ECC is the low socio-economical level of the parents.

Children with ECC have a high frequency of sugar consumption, not only of fluids given in the nursing bottle, but also of sweetened solid foods characteristic is likely to be one of the most

significant caries risk factors. World Health Organization (WHO) recommended that non-milk products and cellular extrinsic sugar should not represent more than 10% of the total daily caloric intake, and that sugars should comprise no more than 10-20% of the human diet. While the scientific basis of the messages to promote breastfeeding for general health are well accepted, prolonged and nocturnal breastfeeding is associated with an increased risk of ECC. However, the evidence for such an association is limited and inconsistent, and is based primarily on cross-sectional studies relying on retrospective recall of infant feeding practices.

ECC is a public health problem and it requires involvement of all health professionals that provide care to children. Oral health cannot be seen as separate from general health. The objective of dentistry surpasses the preservation of the teeth; it aims at maintaining oral and systemic health.

Children receive notions of oral care and values from their families. Therefore, we should change the risk factors for caries development by establishing a partnership with the family, having prevention as the main focus. Prevention is inexpensive, but it demands efforts from family members, who sometimes are not aware of the consequences that caries can bring, or when they realize the problem, the disease is already installed resulting in consequences to the child and family's life.

Essential References

American Academy of Paediatric Dentistry (2008). Definition of Early Childhood Caries (ECC).

http://www.aapd.org/media/policies_guidelines/d_ecc.pdf

Bernabè E, Hobdell MH. Is income inequality related to childhood dental caries in rich countries? JADA 2010;141:143-149.

Drury TF, Horowitz AM, Ismail AI, Maertens MP, Rozier RG, Selwitz RH. Diagnosing and reporting early childhood caries for research purposes. A report of a workshop sponsored by the National Institute of Dental and Craniofacial Research, the Health Resources and Services Administration, and the Health Care Financing Administration. J Public Health Dent 1999;59:192-7.

Feldens CA, Giugliani ER, Vigo Á, Vítolo MR. Early feeding practices and severe early childhood caries in four-year-old children from southern Brazil: a birth cohort study. Caries Res 2010;44:445-52.

<http://www.prisma-statement.org>

Maragakís GM, Kapetanákou DN, Manios Y. Caries prevalence and location and dental treatment needs in preschoolers in Athens--GENESIS project. Community Dent Health 2007;24:264-67.

Niji R, Arita K, Abe Y, Lucas ME, Nishino M, Mitome M. Maternal age at birth and other risk factors in early childhood caries. Pediatr Dent 2010;32:493-98.

Nunn ME, Dietrich T, Singh HK, Henshaw MM, Kressin NR. Prevalence of early childhood caries among very young urban Boston children compared with US children. J Public Health Dent 2009;69:156-62.

Psoter WJ, Zhang H, Pendrys DG, Morse DE, Mayne ST. Classification of dental caries patterns in the primary dentition: a multidimensional scaling analysis. *Community Dent Oral Epidemiol* 2003;31:231-8.

Thitasomakul S, Piwat S, Thearmontree A, Chankanka O, Pithpornchaiyakul W, Madyusoh S. Risks for early childhood caries analyzed by negative binomial models. *J Dent Res* 2009;88:137-41.

Tiberia MJ, Milnes AR, Feigal RJ, Morley KR, Richardson DS, Croft WG, Cheung WS. Risk factors for early childhood caries in Canadian preschool children seeking care. *Pediatr Dent* 2007;29:201-8.

Tyagi R. The prevalence of nursing caries in Davangere preschool children and its relationship with feeding practices and socioeconomic status of the family. *J Indian Soc Pedod Prev Dent* 2008;26:153-7.

Figure 1. Flow chart of the study




Table 1. Papers selected for the review.

<i>Authors</i>	<i>Age group</i>	<i>Size</i>	<i>Country</i>	<i>Prevalence/dmft mean</i>
Thitasomakul <i>et al.</i> (2009)	9-18 months	495	Thailand	Mean range 0.1±0.4 2.8±2.7
Nunn <i>et al.</i> (2009)	1-3 years	787	United States	3.0 %
Barnabé <i>et al.</i> (2010)	5-6 years	--	22 countries	1.68±0.59 mean (0.70-3.31 range)
Niji <i>et al.</i> (2010)	1.5-3 years	646	Japan	41 %
Maragakis <i>et al.</i> (2007)	2-6 years	684	Greece	60.1 %
Tyagi (2008)	2-6 years	813	India	19.2 %
Tiberia <i>et al.</i> (2007)	1-5 years	137	Canada	74.0 %
Feldens <i>et al.</i> (2010)	4 years	340	Brazil	26.0% simple-ECC 37.0 S-ECC

Table 2. Association between background factors (socio-economical levels, dietary behaviours, oral hygiene) and ECC.

<i>Authors</i>	<i>Background factors associated</i>
Thitasomakul <i>et al.</i> (2009)	<ul style="list-style-type: none"> • Children aged 12-18 months • Low income • Mothers did not have a daily intake of milk • No calcium during pregnancy • Children who were breast-fed • Mother's poor oral health status • Mothers had only primary school education • Mothers with 10 or more decayed teeth • Children who were not fed cooked rice or commercial cereal by the age of 3 months • Children who had soft drinks at 9 months • Children who had local traditional desserts at 9 months • Children who had started eating vegetables later than 6 months • Sweetened food • Sugary food by the age of 5 months • Soft drink • Sugary snacks • Children who did not have their teeth brushed at 9 months
Nunn <i>et al.</i> (2009)	<ul style="list-style-type: none"> • Race • History of a child's visit to the dentist • Parent's education • Annual household income
Barnabé <i>et al.</i> (2010)	<ul style="list-style-type: none"> • Gini coefficient
Niji <i>et al.</i> (2010)	<ul style="list-style-type: none"> • Mother's age at childbirth (< 22 years) • Frequency of between meal snacks more than 4 time a day • Child's CAT score of 1.5 year old equal to greater than +1.5
Tyagi (2008)	<ul style="list-style-type: none"> • Children who were bottle-fed • Use of dummy
Tiberia <i>et al.</i> (2007)	<ul style="list-style-type: none"> • Leaving the bottle with the child • Having problems brushing • Holding liquids in the mouth for prolonged time • Being Caucasian
Feldens <i>et al.</i> (2010)	<ul style="list-style-type: none"> • Breastfeeding duration • Frequency of breastfeeding • Night-time bottle use for liquids other than milk • High density of sugar

-
- High density of lipids
 - Maternal schooling
 - Per capita income
 - Teeth at 12 months
-

Appendix 1. List of 411 paper retrieved by MESH search. Papers are listed in date of publishing order.

1. Erdem AP, Guven Y, Balli B, Ilhan B, Sepet E, Ulukapi I, Aktoren O. Success rates of mineral trioxide aggregate, ferric sulfate, and formocresol pulpotomies: a 24-month study. *Pediatr Dent* 2011;33:165-70.
2. Buzalaf MA, Levy SM. Fluoride intake of children: considerations for dental caries and dental fluorosis. *Monogr Oral Sci* 2011;22:1-19.
3. Foster GR, Downer MC, Tickle M. Modelling the impact of process variables in community fluoridated milk schemes on a population of UK schoolchildren. *Br Dent J* 2011 May 28;210:E17.
4. Rayen R, Hariharan VS, Elavazhagan N, Kamalendran N, Varadarajan R. Dental management of hemophiliac child under general anesthesia. *J Indian Soc Pedod Prev Dent* 2011;29:74-9.
5. Lukacs JR. Gender differences in oral health in South Asia: metadata imply multifactorial biological and cultural causes. *Am J Hum Biol* 2011;23:398-411.
6. Yengopal V, Bhayat A, Coogan M. Pediatric oral HIV research in the developing world. *Adv Dent Res* 2011;23:61-6.
7. Frazão P. Effectiveness of the bucco-lingual technique within a school-based supervised toothbrushing program on preventing caries: a randomized controlled trial. *BMC Oral Health* 2011;22:11:11.
8. Curtis B, Warren E, Pollicino C, Evans RW, Schwarz E, Sbaraini A. The Monitor Practice Programme: is non-invasive management of dental caries in private practice cost-effective? *Aust Dent J* 2011;;56:48-55.
9. Maupomé G, Karanja N, Ritenbaugh C, Lutz T, Aickin M, Becker T. Dental caries in American Indian toddlers after a community-based beverage intervention. *Ethn Dis* 2010;20:444-50.
10. Wong MC, Clarkson J, Glenny AM, Lo EC, Marinho VC, Tsang BW, Walsh T, Worthington HV. Cochrane reviews on the benefits/risks of fluoride toothpastes. *J Dent Res* 2011;90:573-9.
11. Arora A, Scott JA, Bhole S, Do L, Schwarz E, Blinkhorn AS. Early childhood feeding practices and dental caries in preschool children: a multi-centre birth cohort study. *BMC Public Health* 2011;12:11:28.
12. Cury JA, Oliveira MJ, Martins CC, Tenuta LM, Paiva SM. Available fluoride in toothpastes used by Brazilian children. *Braz Dent J* 2010;21:396-400.
13. Taji S, Seow WK. A literature review of dental erosion in children. *Aust Dent J* 2010;55:358-67.
14. Ferracane J, Hilton T, Korpak A, Gillette J, McIntyre PS, Berg J; Northwest PRECEDENT. Use of caries prevention services in the Northwest PRECEDENT dental network. *Community Dent Oral Epidemiol* 2011;39:69-78.
15. Peters MC, Tallman JA, Braun TM, Jacobson JJ. Clinical reduction of *S. mutans* in pre-school children using a novel liquorice root extract lollipop: a pilot study. *Eur Arch Paediatr Dent* 2010;11:274-8.
16. Martignon S, Tellez M, Santamaría RM, Gomez J, Ekstrand KR. Sealing distal proximal caries lesions in first primary molars: efficacy after 2.5 years. *Caries Res* 2010;44:562-70.
17. van der Zee V, van Amerongen WE. Short communication: Influence of preformed metal crowns (Hall technique) on the occlusal vertical dimension in the primary dentition. *Eur Arch Paediatr Dent* 2010;11:225-7.
18. Armfield JM. Community effectiveness of public water fluoridation in reducing children's dental disease. *Public Health Rep* 2010;125:655-64.
19. Vilhena FV, Olympio KP, Lauris JR, Delbem AC, Buzalaf MA. Low-fluoride acidic dentifrice: a randomized clinical trial in a fluoridated area. *Caries Res* 2010;44:478-84.
20. Novaes TF, Matos R, Raggio DP, Imperato JC, Braga MM, Mendes FM. Influence of the discomfort reported by children on the performance of approximal caries detection methods. *Caries Res* 2010;44:465-71.

Dr.ssa Giovanna Congiu EARLY CHILDHOOD CARIES (ECC) IS STILL A PROBLEM? AN EPIDEMIOLOGICAL AND AETIOLOGICAL APPROACH.

Università degli Studi di Sassari

21. Parker EJ, Jamieson LM, Broughton J, Albino J, Lawrence HP, Roberts-Thomson K. The oral health of Indigenous children: a review of four nations. *J Paediatr Child Health* 2010;46:483-6.
22. Petti S. Why guidelines for early childhood caries prevention could be ineffective amongst children at high risk. *J Dent* 2010;38:946-55.
23. Orhan AI, Oz FT, Orhan K. Pulp exposure occurrence and outcomes after 1- or 2-visit indirect pulp therapy vs complete caries removal in primary and permanent molars. *Pediatr Dent* 2010;32:347-55.
24. Isik EE, Olmez A, Akca G, Sultan N. A microbiological assessment of polymer and conventional carbide burs in caries removal. *Pediatr Dent* 2010;32:316-23.
25. Wang SJ, Briskie D, Hu JC, Majewski R, Inglehart MR. Illustrated information for parent education: parent and patient responses. *Pediatr Dent* 2010;32:295-303.
26. Aminabadi NA, Farahani RM, Oskouei SG. Formocresol versus calcium hydroxide direct pulp capping of human primary molars: two year follow-up. *J Clin Pediatr Dent* 2010;34:317-21.
27. Braga MM, Martignon S, Ekstrand KR, Ricketts DN, Imparato JC, Mendes FM. Parameters associated with active caries lesions assessed by two different visual scoring systems on occlusal surfaces of primary molars - a multilevel approach. *Community Dent Oral Epidemiol* 2010;38:549-58.
28. Roberts-Thomson KF, Slade GD, Bailie RS, Endean C, Simmons B, Leach AJ, Raye I, Morris PS. A comprehensive approach to health promotion for the reduction of dental caries in remote Indigenous Australian children: a clustered randomised controlled trial. *Int Dent J* 2010;60:245-9.
29. Lawrence HP. Oral health interventions among Indigenous populations in Canada. *Int Dent J* 2010;60:229-34.
30. Albino JE, Orlando VA. Promising directions for caries prevention with American Indian and Alaska Native children. *Int Dent J* 2010;60:216-22.
31. Slade GD, Bailie RS, Roberts-Thomson K, Leach AJ, Raye I, Endean C, Simmons B, Morris P. Effect of health promotion and fluoride varnish on dental caries among Australian Aboriginal children: results from a community-randomized controlled trial. *Community Dent Oral Epidemiol* 2011 Feb;39:29-43.
32. Qvist V, Poulsen A, Teglers PT, Mjör IA. Fluorides leaching from restorative materials and the effect on adjacent teeth. *Int Dent J* 2010;60:156-60.
33. Carvalho DM, Salazar M, Oliveira BH, Coutinho ES. Fluoride varnishes and decrease in caries incidence in preschool children: a systematic review. *Rev Bras Epidemiol* 2010;13:139-49.
34. Allen G, Logan R, Gue S. Oral manifestations of cancer treatment in children: a review of the literature. *Clin J Oncol Nurs* 2010;14:481-90.
35. Strippel H. Effectiveness of structured comprehensive paediatric oral health education for parents of children less than two years of age in Germany. *Community Dent Health* 2010;27:74-80.
36. Deepa G, Shobha T. A clinical evaluation of two glass ionomer cements in primary molars using atraumatic restorative treatment technique in India: 1 year follow up. *Int J Paediatr Dent* 2010;20:410-8.
37. Aminabadi NA, Ghoreishizadeh A, Ghoreishizadeh M, Oskouei SG. Can drawing be considered a projective measure for children's distress in paediatric dentistry? *Int J Paediatr Dent* 2011;21:1-12.
38. James P, Parnell C, Whelton H. The caries-preventive effect of chlorhexidine varnish in children and adolescents: a systematic review. *Caries Res* 2010;44:333-40.
39. Simratvir M, Singh N, Chopra S, Thomas AM. Efficacy of 10% Povidone Iodine in children affected with early childhood caries: an in vivo study. *J Clin Pediatr Dent* 2010;34:233-8.
40. Warren E, Pollicino C, Curtis B, Evans W, Sbaraini A, Schwarz E. Modeling the long-term cost-effectiveness of the caries management system in an Australian population. *Value Health* 2010;13:750-60.
41. Plutzer K, Keirse MJ. Incidence and prevention of early childhood caries in one- and two-parent families. *Child Care Health Dev* 2011;37:5-10.
42. Simancas-Pallares MA, Díaz-Caballero AJ, Luna-Ricardo LM. Mineral trioxide aggregate in primary teeth pulpotomy. A systematic literature review. *Med Oral Patol Oral Cir Bucal* 2010;15:e942-6.
43. Ansari G, Ranjpour M. Mineral trioxide aggregate and formocresol pulpotomy of primary teeth: a 2-year follow-up. *Int Endod J* 2010;43:413-8.
44. Mansour Ockell N, Bågesund M. Reasons for extractions, and treatment preceding caries-related extractions in 3-8 year-old children. *Eur Arch Paediatr Dent* 2010;11:122-30.

45. Marks LA, Faict N, Welbury RR. Literature review: Restorations of class II cavities in the primary dentition with compomers. *Eur Arch Paediatr Dent* 2010;11:109-14.
46. Parisotto TM, Steiner-Oliveira C, Silva CM, Rodrigues LK, Nobre-dos-Santos M. Early childhood caries and mutans streptococci: a systematic review. *Oral Health Prev Dent* 2010;8:59-70.
47. Doherty MA, Blinkhorn AS, Vane ES. Oral health in the Pacific Islands. *Int Dent J* 2010;60:122-8.
48. Harrison R, Veronneau J, Leroux B. Design and implementation of a dental caries prevention trial in remote Canadian Aboriginal communities. *Trials* 2010;13:11:54.
49. Sonmez D, Duruturk L. Success rate of calcium hydroxide pulpotomy in primary molars restored with amalgam and stainless steel crowns. *Br Dent J* 2010;8;208:E18.
50. Casagrande L, Bento LW, Dalpian DM, García-Godoy F, de Araujo FB. Indirect pulp treatment in primary teeth: 4-year results. *Am J Dent* 2010;23:34-8.
51. Chandna P, Adlakha VK. Oral health in children guidelines for pediatricians. *Indian Pediatr* 2010;47:323-7.
52. Feldens CA, Giugliani ER, Duncan BB, Drachler Mde L, Vítolo MR. Long-term effectiveness of a nutritional program in reducing early childhood caries: a randomized trial. *Community Dent Oral Epidemiol* 2010;38:324-32.
53. Cunnion DT, Spiro A 3rd, Jones JA, Rich SE, Papageorgiou CP, Tate A, Casamassimo P, Hayes C, Garcia RI. Pediatric oral health-related quality of life improvement after treatment of early childhood caries: a prospective multisite study. *J Dent Child* 2010;77:4-11.
54. Davis EE, Deinard AS, Maiga EW. Doctor, my tooth hurts: the costs of incomplete dental care in the emergency room. *J Public Health Dent* 2010;70:205-10.
55. de Menezes Abreu DM, Leal SC, Frencken JE. Self-report of pain in children treated according to the atraumatic restorative treatment and the conventional restorative treatment--a pilot study. *J Clin Pediatr Dent* 2009;34:151-5.
56. Kotb RM, Abdella AA, El Kateb MA, Ahmed AM. Clinical evaluation of Papacarie in primary teeth. *J Clin Pediatr Dent* 2009;34:117-23.
57. Alves dos Santos MP, Luiz RR, Maia LC. Randomised trial of resin-based restorations in Class I and Class II beveled preparations in primary molars: 48-month results. *J Dent*. 2010;38:451-9.
58. Cleaton-Jones P, Fatti P. Dental caries in children in South Africa and Swaziland: a systematic review 1919-2007. *Int Dent J* 2009;59:363-8.
59. Ertugrul F, Cogulu D, Ozdemir Y, Ersin N. Comparison of conventional versus colored compomers for class II restorations in primary molars: a 12-month clinical study. *Med Princ Pract* 2010;19:148-52.
60. van Gemert-Schriks MC, van Amerongen EW, Aartman IH, Wennink JM, Ten Cate JM, de Soet JJ. The influence of dental caries on body growth in prepubertal children. *Clin Oral Investig* 2011;15:141-9.
61. Wong MC, Glenny AM, Tsang BW, Lo EC, Worthington HV, Marinho VC. Topical fluoride as a cause of dental fluorosis in children. *Cochrane Database Syst Rev* 2010 ;20:CD007693.
62. Ekstrand KR, Bakhshandeh A, Martignon S. Treatment of proximal superficial caries lesions on primary molar teeth with resin infiltration and fluoride varnish versus fluoride varnish only: efficacy after 1 year. *Caries Res* 2010;44:41-6.
63. Freudenthal JJ, Bowen DM. Motivational interviewing to decrease parental risk-related behaviors for early childhood caries. *J Dent Hyg* 2010;84:29-34.
64. Castilho SD, Rocha MA. Pacifier habit: history and multidisciplinary view. *J Pediatr (Rio J)* 2009;85:480-9.
65. Weinstein P, Spiekerman C, Milgrom P. Randomized equivalence trial of intensive and semiannual applications of fluoride varnish in the primary dentition. *Caries Res* 2009;43:484-90.
66. Nakai Y, Shinga-Ishihara C, Kaji M, Moriya K, Murakami-Yamanaka K, Takimura M. Xylitol gum and maternal transmission of mutans streptococci. *J Dent Res* 2010;89:56-60.
67. Mobley C, Marshall TA, Milgrom P, Coldwell SE. The contribution of dietary factors to dental caries and disparities in caries. *Acad Pediatr* 2009;9:410-4.
68. Huston J, Wood AJ. Sharing early preventive oral health with medical colleagues: a dental pain prevention strategy. *J Calif Dent Assoc* 2009;37:723-34.
69. Jaakkola E, Laine CM, Mäyränpää MK, Falck A, Ignatius J, Mäkitie O. Calvarial doughnut lesions and osteoporosis: a new three-generation family and review. *Am J Med Genet A* 2009;149:2371-7.
70. Milgrom P, Zero DT, Tanzer JM. An examination of the advances in science and technology of prevention of tooth decay in young children since the Surgeon General's Report on Oral Health. *Acad Pediatr* 2009;9:404-9.

71. Ellwood RP, Cury JA. How much toothpaste should a child under the age of 6 years use? *Eur Arch Paediatr Dent* 2009;10:168-74.
72. Twetman S. Caries prevention with fluoride toothpaste in children: an update. *Eur Arch Paediatr Dent* 2009;10:162-7.
73. Jyoti S, Shashikiran ND, Reddy VV. Effect of lactoperoxidase system containing toothpaste on cariogenic bacteria in children with early childhood caries. *J Clin Pediatr Dent* 2009;33:299-303.
74. Söderling EM. Xylitol, mutans streptococci, and dental plaque. *Adv Dent Res* 2009;21:74-8.
75. Kramer MS, Matush L, Bogdanovich N, Aboud F, Mazer B, Fombonne E, Collet JP, Hodnett E, Mironova E, Igumnov S, Chalmers B, Dahhou M, Platt RW. Health and development outcomes in 6.5-y-old children breastfed exclusively for 3 or 6 mo. *Am J Clin Nutr* 2009;90:1070-4.
76. Stecksén-Blicks C, Sjöström I, Twetman S. Effect of long-term consumption of milk supplemented with probiotic lactobacilli and fluoride on dental caries and general health in preschool children: a cluster-randomized study. *Caries Res* 2009;43:374-81.
77. Do LG, Spencer AJ, Ha DH. Association between dental caries and fluorosis among South Australian children. *Caries Res* 2009;43:366-73.
78. Lasso EM, Tavares MC, Silva JY, Urban Cde A. Severe early childhood caries: an integral approach. *J Pediatr (Rio J)* 2009;85:295-300.
79. Lula EC, Monteiro-Neto V, Alves CM, Ribeiro CC. Microbiological analysis after complete or partial removal of carious dentin in primary teeth: a randomized clinical trial. *Caries Res* 2009;43:354-8.
80. Yee R, Holmgren C, Mulder J, Lama D, Walker D, van Palenstein Helder W. Efficacy of silver diamine fluoride for Arresting Caries Treatment. *J Dent Res* 2009;88:644-7.
81. Sakai VT, Moretti AB, Oliveira TM, Fornetti AP, Santos CF, Machado MA, Abdo RC. Pulpotomy of human primary molars with MTA and Portland cement: a randomised controlled trial. *Br Dent J* 2009;207:E5.
82. Casagrande L, Falster CA, Di Hipolito V, De Góes MF, Straffon LH, Nör JE, de Araujo FB. Effect of adhesive restorations over incomplete dentin caries removal: 5-year follow-up study in primary teeth. *J Dent Child (Chic)* 2009;76:117-22.
83. dos Santos Pinheiro R, França TT, Ribeiro CM, Leão JC, de Souza IP, Castro GF. Oral manifestations in human immunodeficiency virus infected children in highly active antiretroviral therapy era. *J Oral Pathol Med* 2009;38:613-22.
84. Klaassen MA, Veerkamp JS, Hoogstraten J. Young children's Oral Health-Related Quality of Life and dental fear after treatment under general anaesthesia: a randomized controlled trial. *Eur J Oral Sci* 2009;117:273-8.
85. Milgrom P, Ly KA, Tut OK, Mancil L, Roberts MC, Briand K, Gancio MJ. Xylitol pediatric topical oral syrup to prevent dental caries: a double-blind randomized clinical trial of efficacy. *Arch Pediatr Adolesc Med* 2009;163:601-7.
86. Fontana M, Catt D, Eckert GJ, Ofner S, Toro M, Gregory RL, Zandona AF, Eggertsson H, Jackson R, Chin J, Zero D, Sissons CH. Xylitol: effects on the acquisition of cariogenic species in infants. *Pediatr Dent* 2009;31:257-66.
87. Duque C, Negrini Tde C, Sacono NT, Spolidorio DM, de Souza Costa CA, Hebling J. Clinical and microbiological performance of resin-modified glass-ionomer liners after incomplete dentine caries removal. *Clin Oral Investig* 2009;13:465-71.
88. Meyer K, Geurtsen W, Günay H. An early oral health care program starting during pregnancy: results of a prospective clinical long-term study. *Clin Oral Investig* 2010;14:257-64.
89. Twetman S, Fontana M. Patient caries risk assessment. *Monogr Oral Sci* 2009;21:91-101.
90. Milgrom P, Tut OK. Evaluation of Pacific Islands Early Childhood Caries Prevention Project: Republic of the Marshall Islands. *J Public Health Dent* 2009;69:201-3.
91. Cruz SS, Costa Mda C, Gomes-Filho IS, Barreto ML, dos Santos CA, Martins AG, Passos Jde S, de Freitas CO, Sampaio FP, Cerqueira Ede M. Periodontal therapy for pregnant women and cases of low birthweight: an intervention study. *Pediatr Int* 2010;52:57-64.
92. Yengopal V, Harneker SY, Patel N, Siegfried N. Dental fillings for the treatment of caries in the primary dentition. *Cochrane Database Syst Rev* 2009;15:CD004483.
93. Barja-Fidalgo F, Maroun S, de Oliveira BH. Effectiveness of a glass ionomer cement used as a pit and fissure sealant in recently erupted permanent first molars. *J Dent Child (Chic)* 2009;76:34-40.
94. Mohebbi SZ, Virtanen JI, Vahid-Golpayegani M, Vehkalahti MM. A cluster randomised trial of effectiveness of educational intervention in primary health care on early childhood caries. *Caries Res* 2009;43:110-8.

95. Meurman P, Pienihäkkinen K, Eriksson AL, Alanen P. Oral health programme for preschool children: a prospective, controlled study. *Int J Paediatr Dent* 2009;19:263-73.
96. Bagramian RA, Garcia-Godoy F, Volpe AR. The global increase in dental caries. A pending public health crisis. *Am J Dent* 2009;22:3-8.
97. Garrocho-Rangel A, Flores H, Silva-Herzog D, Hernandez-Sierra F, Mandeville P, Pozos-Guillen AJ. Efficacy of EMD versus calcium hydroxide in direct pulp capping of primary molars: a randomized controlled clinical trial. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009;107:733-8.
98. Nunn ME, Dietrich T, Singh HK, Henshaw MM, Kressin NR. Prevalence of early childhood caries among very young urban Boston children compared with US children. *J Public Health Dent* 2009;69:156-62.
99. dos Santos MP, Passos M, Luiz RR, Maia LC. A randomized trial of resin-based restorations in class I and class II beveled preparations in primary molars: 24-month results. *J Am Dent Assoc* 2009;140:156-66.
100. Williamson R, Oueis H, Casamassimo PS, Thikkurissy S. Association between early childhood caries and behavior as measured by the Child Behavior Checklist. *Pediatr Dent* 2008;30:505-9.
101. Andruskeviciene V, Milciuviene S, Bendoraitiene E, Saldunaite K, Vasiliauskiene I, Slabsinskiene E, Narbutaite J. Oral health status and effectiveness of caries prevention programme in kindergartens in Kaunas city (Lithuania). *Oral Health Prev Dent* 2008;6:343-8.
102. Chu CH, Lo EC. Promoting caries arrest in children with silver diamine fluoride: a review. *Oral Health Prev Dent* 2008;6:315-21.
103. Plutzer K, Spencer AJ. Efficacy of an oral health promotion intervention in the prevention of early childhood caries. *Community Dent Oral Epidemiol* 2008;36:335-46.
104. Kagihara LE, Niederhauser VP, Stark M. Assessment, management, and prevention of early childhood caries. *J Am Acad Nurse Pract* 2009;21:1-10.
105. Poureslami HR, Van Amerongen WE. Early Childhood Caries (ECC): an infectious transmissible oral disease. *Indian J Pediatr* 2009;76:191-4.
106. Kindelan SA, Day P, Nichol R, Willmott N, Fayle SA; British Society of Paediatric Dentistry. UK National Clinical Guidelines in Paediatric Dentistry: stainless steel preformed crowns for primary molars. *Int J Paediatr Dent* 2008;18:20-8.
107. Carrillo CM, Tanaka MH, Cesar MF, Camargo MA, Juliano Y, Novo NF. Use of papain gel in disabled patients. *J Dent Child (Chic)* 2008;75:222-8.
108. Hauser-Gerspach I, Pfäffli-Savtchenko V, Dähnhardt JE, Meyer J, Lussi A. Comparison of the immediate effects of gaseous ozone and chlorhexidine gel on bacteria in cavitated carious lesions in children in vivo. *Clin Oral Investig* 2009;13:287-91.
109. Section on Pediatric Dentistry and Oral Health. Preventive oral health intervention for pediatricians. *Pediatrics* 2008;122:1387-94.
110. Grewal N, Seth R. Comparative in vivo evaluation of restoring severely mutilated primary anterior teeth with biological post and crown preparation and reinforced composite restoration. *J Indian Soc Pedod Prev Dent* 2008;26:141-8.
111. Ismail AI, Hasson H. Fluoride supplements, dental caries and fluorosis: a systematic review. *J Am Dent Assoc* 2008;139:1457-68.
112. Lobo PL, de Carvalho CB, Fonseca SG, de Castro RS, Monteiro AJ, Fonteles MC, Fonteles CS. Sodium fluoride and chlorhexidine effect in the inhibition of mutans streptococci in children with dental caries: a randomized, double-blind clinical trial. *Oral Microbiol Immunol* 2008;23:486-91.
113. Hosoya Y, Taguchi T, Arita S, Tay FR. Clinical evaluation of polypropylene glycol-based caries detecting dyes for primary and permanent carious dentin. *J Dent* 2008;36:1041-7.
114. Mohamed N. A comparison of two liner materials for use in the ferric sulfate pulpotomy. *SADJ* 2008;63:338, 340-2.
115. Vadiakas G. Case definition, aetiology and risk assessment of early childhood caries (ECC): a revisited review. *Eur Arch Paediatr Dent* 2008;9:114-25.
116. Alfaro EV, Aps JK, Martens LC. Oral implications in children with gastroesophageal reflux disease. *Curr Opin Pediatr* 2008;20:576-83.
117. Miller EK, Vann WF Jr. The use of fluoride varnish in children: a critical review with treatment recommendations. *J Clin Pediatr Dent* 2008;32:259-64.

118. Franco S, Theriot J, Greenwell A. The influence of early counselling on weaning from a bottle. *Community Dent Health* 2008;25:115-8.
119. Oliveira CA, Dias PF, Dos Santos MP, Maia LC. Split mouth randomized controlled clinical trial of beveled cavity preparations in primary molars: an 18-Month follow up. *J Dent* 2008;36:754-8.
120. Marja-Leena M, Paivi R, Sirkka J, Ansa O, Matti S. Childhood caries is still in force: a 15-year follow-up. *Acta Odontol Scand* 2008;66:189-92.
121. Orhan AI, Oz FT, Ozcelik B, Orhan K. A clinical and microbiological comparative study of deep carious lesion treatment in deciduous and young permanent molars. *Clin Oral Investig*. 2008;12:369-78.
122. Roberts MW. Dental health of children: where we are today and remaining challenges. *J Clin Pediatr Dent* 2008;32:231-4.
123. Aminabadi NA, Farahani RM, Gajan EB. A clinical study of formocresol pulpotomy versus root canal therapy of vital primary incisors. *J Clin Pediatr Dent* 2008;32:211-4.
124. Versloot J, Veerkamp JS, Hoogstraten J. Pain behaviour and distress in children during two sequential dental visits: comparing a computerised anaesthesia delivery system and a traditional syringe. *Br Dent J* 2008;12:205:E2.
125. Vilan Xavier AC, Pinto Silva LC, Oliveira P, Villamarim Soares R, de Almeida Cruz R. A review and dental management of persons with craniosynostosis anomalies. *Spec Care Dentist* 2008;28:96-100.
126. de Amorim RG, Leal SC, Bezerra AC, de Amorim FP, de Toledo OA. Association of chlorhexidine and fluoride for plaque control and white spot lesion remineralization in primary dentition. *Int J Paediatr Dent* 2008;18:446-51.
127. Moretti AB, Sakai VT, Oliveira TM, Fornetti AP, Santos CF, Machado MA, Abdo RC. The effectiveness of mineral trioxide aggregate, calcium hydroxide and formocresol for pulpotomies in primary teeth. *Int Endod J* 2008;41:547-55.
128. Whittle JG, Whitehead HF, Bishop CM. A randomised control trial of oral health education provided by a health visitor to parents of pre-school children. *Community Dent Health* 2008;25:28-32.
129. Noorollahian H. Comparison of mineral trioxide aggregate and formocresol as pulp medicaments for pulpotomies in primary molars. *Br Dent J* 2008;14:204:E20.
130. Lawrence HP, Binguis D, Douglas J, McKeown L, Switzer B, Figueiredo R, Laporte A. A 2-year community-randomized controlled trial of fluoride varnish to prevent early childhood caries in Aboriginal children. *Community Dent Oral Epidemiol* 2008;36:503-16.
131. Zurn D, Seale NS. Light-cured calcium hydroxide vs formocresol in human primary molar pulpotomies: a randomized controlled trial. *Pediatr Dent* 2008;30:34-41.
132. van Gemert-Schriks MC, van Amerongen WE, ten Cate JM, Aartman IH. The effect of different dental treatment strategies on the oral health of children: a longitudinal randomised controlled trial. *Clin Oral Investig* 2008;12:361-8.
133. Cerqueira DF, Mello-Moura AC, Santos EM, Guedes-Pinto AC. Cytotoxicity, histopathological, microbiological and clinical aspects of an endodontic iodoform-based paste used in pediatric dentistry: a review. *J Clin Pediatr Dent* 2008;32:105-10.
134. Chu CH, Lo EC. Microhardness of dentine in primary teeth after topical fluoride applications. *J Dent* 2008;36:387-91.
135. Twetman S. Prevention of early childhood caries (ECC)--review of literature published 1998-2007. *Eur Arch Paediatr Dent* 2008;9:12-8.
136. Atieh M. Stainless steel crown versus modified open-sandwich restorations for primary molars: a 2-year randomized clinical trial. *Int J Paediatr Dent* 2008;18:325-32.
137. Azarpazhooh A, Main PA. Fluoride varnish in the prevention of dental caries in children and adolescents: a systematic review. *J Can Dent Assoc* 2008;74:73-9.
138. Prabhakar AR, Raju OS, Kurthukoti AJ, Satish V. Evaluation of the clinical behavior of resin modified glass ionomer cement on primary molars: a comparative one-year study. *J Contemp Dent Pract* 2008;9:130-7.
139. Maragakis GM, Kapetanidou DN, Manios Y. Caries prevalence and location and dental treatment needs in preschoolers in Athens--GENESIS project. *Community Dent Health* 2007;24:264-7.
140. Krause F, Braun A, Lotz G, Kneist S, Jepsen S, Eberhard J. Evaluation of selective caries removal in deciduous teeth by a fluorescence feedback-controlled Er:YAG laser in vivo. *Clin Oral Investig* 2008;12:209-15.
141. Lanigan J, Turnbull B, Singhal A. Toddler diets in the UK: deficiencies and imbalances. 2. Relationship of toddler diet to later health. *J Fam Health Care* 2007;17:197-200.

142. Quiñonez RB, Pahel BT, Rozier RG, Stearns SC. Follow-up preventive dental visits for Medicaid-enrolled children in the medical office. *J Public Health Dent* 2008;63:131-8.
143. Southward LH, Robertson A, Edelstein BL, Hanna H, Wells-Parker E, Baggett DH, Eklund NP, Crall JJ, Silberman SL, Parrish DR. Oral health of young children in Mississippi Delta child care centers: a second look at early childhood caries risk assessment. *J Public Health Dent* 2008;68:188-95.
144. Casagrande L, Bento LW, Rerim SO, Lucas Ede R, Dalpian DM, de Araujo FB. In vivo outcomes of indirect pulp treatment using a self-etching primer versus calcium hydroxide over the demineralized dentin in primary molars. *J Clin Pediatr Dent* 2008;33:131-5.
145. Fan X, Li X, Wan H, Hu D, Zhang YP, Volpe AR, DeVizio W. Clinical investigation of the anticaries efficacy of a 1.14% sodium monofluorophosphate (SMFP) calcium carbonate-based dentifrice: a two-year caries clinical trial on children in China. *J Clin Dent* 2008;19:134-7.
146. Yost J, Li Y. Promoting oral health from birth through childhood: prevention of early childhood caries. *MCN Am J Matern Child Nurs* 2008;33:17-23.
147. Misra S, Tahmassebi JF, Brosnan M. Early childhood caries--a review. *Dent Update* 2007;34:556-8, 561-2, 564.
148. Young DA, Featherstone JD, Roth JR, Anderson M, Autio-Gold J, Christensen GJ, Fontana M, Kutsch VK, Peters MC, Simonsen RJ, Wolff MS. Caries management by risk assessment: implementation guidelines. *J Calif Dent Assoc* 2007;35:799-805.
149. Bánóczy J, Rugg-Gunn AJ. Caries prevention through the fluoridation of milk. A review. *Fogorv Sz* 2007;100:185-192, 177-84.
150. Lima TJ, Ribeiro CC, Tenuta LM, Cury JA. Low-fluoride dentifrice and caries lesion control in children with different caries experience: a randomized clinical trial. *Caries Res* 2008;42:46-50.
151. Tuna D, Olmez A. Clinical long-term evaluation of MTA as a direct pulp capping material in primary teeth. *Int Endod J* 2008;41:273-8.
152. Harrison R, Benton T, Everson-Stewart S, Weinstein P. Effect of motivational interviewing on rates of early childhood caries: a randomized trial. *Pediatr Dent* 2007;29:16-22.
153. Lam DK, Sándor GK, Holmes HI, Carmichael RP, Clokie CM. Marble bone disease: a review of osteopetrosis and its oral health implications for dentists. *J Can Dent Assoc* 2007;73:839-43.
154. Leskinen K, Salo S, Suni J, Larmas M. Comparison of dental health in sealed and non-sealed first permanent molars: 7 years follow-up in practice-based dentistry. *J Dent* 2008;36:27-32.
155. Jeevarathan J, Deepthi A, Muthu MS, Rathna Prabhu V, Chamundeeswari GS. Effect of fluoride varnish on *Streptococcus mutans* counts in plaque of caries-free children using Dentocult SM strip mutans test: a randomized controlled triple blind study. *J Indian Soc Pedod Prev Dent* 2007;25:157-63.
156. Turnbull B, Lanigan J, Singhal A. Toddler diets in the U.K.: deficiencies and imbalances. 1. Risk of micronutrient deficiencies. *J Fam Health Care* 2007;17:167-70.
157. Slade GD, Rozier RG, Zeldin LP, Margolis PA. Training pediatric health care providers in prevention of dental decay: results from a randomized controlled trial. *BMC Health Serv Res* 2007;7:176.
158. Cinar DN. The advantages and disadvantages of pacifier use. *Contemp Nurse* 2004;17:109-12.
159. Ammari JB, Baqain ZH, Ashley PF. Effects of programs for prevention of early childhood caries. A systematic review. *Med Princ Pract* 2007;16:437-42.
160. Mornet E. Hypophosphatasia. *Orphanet J Rare Dis* 2007;2:40.
161. Büyükgöral B, Cehreli ZC. Effect of different adhesive protocols vs calcium hydroxide on primary tooth pulp with different remaining dentin thicknesses:24-month results. *Clin Oral Investig* 2008;12:91-6.
162. Galganny-Almeida A, Queiroz MC, Leite AJ. The effectiveness of a novel infant tooth wipe in high caries-risk babies 8 to 15 months old. *Pediatr Dent* 2007;29:337-42.
163. Pine CM, Curnow MM, Burnside G, Nicholson JA, Roberts AJ. Caries prevalence four years after the end of a randomised controlled trial. *Caries Res* 2007;41:431-6.
164. Naidoo S, Myburgh N. Nutrition, oral health and the young child. *Matern Child Nutr* 2007;3:312-21.
165. Bowley NA, Pentz-Kluyts MA, Bourne LT, Marino LV. Feeding the 1 to 7-year-old child. A support paper for the South African paediatric food-based dietary guidelines. *Matern Child Nutr* 2007;3:281-91.
166. Nield LS, Stenger JP, Kamat D. Common pediatric dental dilemmas. *Clin Pediatr (Phila)* 2008;47:99-105.

167. Tiberia MJ, Milnes AR, Feigal RJ, Morley KR, Richardson DS, Croft WG, Cheung WS. Risk factors for early childhood caries in Canadian preschool children seeking care. *Pediatr Dent*. 2007;29:201-8.
168. Law V, Seow WK, Townsend G. Factors influencing oral colonization of mutans streptococci in young children. *Aust Dent J* 2007;52:93-100.
169. Franzon R, Casagrande L, Pinto AS, García-Godoy F, Maltz M, de Araujo FB. Clinical and radiographic evaluation of indirect pulp treatment in primary molars: 36 months follow-up. *Am J Dent* 2007;20:189-92.
170. Karjalainen S. Eating patterns, diet and dental caries. *Dent Update* 2007;34:295-8, 300.
171. Davies GM, Duxbury JT, Boothman NJ, Davies RM. Challenges associated with the evaluation of a dental health promotion programme in a deprived urban area. *Community Dent Health* 2007;24:117-21.
172. Feldens CA, Vítolo MR, Drachler Mde L. A randomized trial of the effectiveness of home visits in preventing early childhood caries. *Community Dent Oral Epidemiol* 2007;35:215-23.
173. Blignaut E. Oral health needs of HIV/AIDS orphans in Gauteng, South Africa. *AIDS Care* 2007;19:532-8.
174. Krämer N, Lohbauer U, Frankenberger R. Restorative materials in the primary dentition of poli-carious patients. *Eur Arch Paediatr Dent* 2007 Mar;8:29-35.
175. Javier JR, Huffman LC, Mendoza FS. Filipino child health in the United States: do health and health care disparities exist? *Prev Chronic Dis* 2007;4:A36.
176. Martignon S, González MC, Santamaría RM, Jácome-Liévano S, Muñoz Y, Moreno P. Oral-health workshop targeted at 0-5-yr. old deprived children's parents and caregivers: effect on knowledge and practices. *J Clin Pediatr Dent* 2006;31:104-8.
177. Marchi JJ, de Araujo FB, Fröner AM, Straffon LH, Nör JE. Indirect pulp capping in the primary dentition: a 4 year follow-up study. *J Clin Pediatr Dent* 2006;31:68-71.
178. Sampaio AL, Pinheiro TG, Furtado PL, Araújo MF, Oliveira CA. Evaluation of early postoperative morbidity in pediatric tonsillectomy with the use of sucralfate. *Int J Pediatr Otorhinolaryngol* 2007;71:645-51.
179. Vargas KG, Packham B, Lowman D. Preliminary evaluation of sodium hypochlorite for pulpotomies in primary molars. *Pediatr Dent* 2006;28:511-7.
180. Selwitz RH, Ismail AI, Pitts NB. Dental caries. *Lancet* 2007;369:51-9.
181. Brennan MT, Kent ML, Fox PC, Norton HJ, Lockhart PB. The impact of oral disease and nonsurgical treatment on bacteremia in children. *J Am Dent Assoc* 2007;138:80-5.
182. Thorild I, Lindau B, Twetman S. Caries in 4-year-old children after maternal chewing of gums containing combinations of xylitol, sorbitol, chlorhexidine and fluoride. *Eur Arch Paediatr Dent* 2006;7:241-5.
183. Elfrink ME, Veerkamp JS, Kalsbeek H. Caries pattern in primary molars in Dutch 5-year-old children. *Eur Arch Paediatr Dent* 2006;7:236-40.
184. Oscarson P, Lif Holgersson P, Sjöström I, Twetman S, Stecksén-Blicks C. Influence of a low xylitol-dose on mutans streptococci colonisation and caries development in preschool children. *Eur Arch Paediatr Dent* 2006;7:142-7.
185. Versloot J, Veerkamp JS, Hoogstraten J. Dental Discomfort Questionnaire for young children following full mouth rehabilitation under general anaesthesia: a follow-up report. *Eur Arch Paediatr Dent* 2006;7:126-9.
186. Foley J. Alternative treatment strategies for carious primary teeth: an overview of the evidence. *Eur Arch Paediatr Dent* 2006;7:73-80.
187. Sheiham A. Dental caries affects body weight, growth and quality of life in pre-school children. *Br Dent J* 2006;201:625-6.
188. Andersson-Wenckert I, Sunnegårdh-Grönberg K. Flowable resin composite as a class II restorative in primary molars: A two-year clinical evaluation. *Acta Odontol Scand* 2006;64:334-40.
189. Saalfield S, Jackson-Allen P. Biopsychosocial consequences of sweetened drink consumption in children 0-6 years of age. *Pediatr Nurs* 2006;32:460-2, 467-71.
190. Spector ND, Kelly SF. Pediatrician's role in screening and treatment: bullying, prediabetes, oral health. *Curr Opin Pediatr* 2006;18:661-70.
191. Ganesh M, Tandon S. Clinical evaluation of FUJI VII sealant material. *J Clin Pediatr Dent* 2006;31:52-7.
192. Wan K, Jing Q, Zhao JZ. Evaluation of oral midazolam as conscious sedation for pediatric patients in oral restoration. *Chin Med Sci J* 2006;21:163-6.

193. Dähnhardt JE, Jaeggi T, Lussi A. Treating open carious lesions in anxious children with ozone. A prospective controlled clinical study. *Am J Dent* 2006;19:267-70.
194. Cleaton-Jones P, Fatti P, Bönecker M. Dental caries trends in 5- to 6-year-old and 11- to 13-year-old children in three UNICEF designated regions--Sub Saharan Africa, Middle East and North Africa, Latin America and Caribbean: 1970-2004. *Int Dent J* 2006;56:294-300.
195. Petti S, Hausen H. Caries-preventive effect of chlorhexidine gel applications among high-risk children. *Caries Res* 2006;40:514-21.
196. Brown LM, Casamassimo PS, Griffen A, Tatakis D. Supragingival calculus in children with gastrostomy feeding: significant reduction with a caregiver-applied tartar-control dentifrice. *Pediatr Dent* 2006;28:410-4.
197. Reeves A, Chiappelli F, Cajulis OS. Evidence-based recommendations for the use of sealants. *J Calif Dent Assoc* 2006;34:540-6.
198. Kirzioglu Z, Gurbuz T, Yilmaz Y. Clinical evaluation of chemomechanical and mechanical caries removal: status of the restorations at 3, 6, 9 and 12 months. *Clin Oral Investig* 2007;11:69-76.
199. Menezes JP, Rosenblatt A, Medeiros E. Clinical evaluation of atraumatic restorations in primary molars: a comparison between 2 glass ionomer cements. *J Dent Child (Chic)* 2006;73:91-7.
200. Thenisch NL, Bachmann LM, Imfeld T, Leisebach Minder T, Steurer J. Are mutans streptococci detected in preschool children a reliable predictive factor for dental caries risk? A systematic review. *Caries Res* 2006;40:366-74.
201. Hosoya Y, Taguchi T, Tay FR. Evaluation of a new caries detecting dye for primary and permanent carious dentin. *J Dent* 2007;35:137-43.
202. Liu JF, Lai YL, Shu WY, Lee SY. Acceptance and efficiency of Er:YAG laser for cavity preparation in children. *Photomed Laser Surg* 2006;24:489-93.
203. Zhan L, Featherstone JD, Gansky SA, Hoover CI, Fujino T, Berkowitz RJ, Den Besten PK. Antibacterial treatment needed for severe early childhood caries. *J Public Health Dent* 2006;66:174-9.
204. Popat H, Dinnage J. Improving cross-cultural awareness. A review of Australian indigenous health for UK dentists. *Br Dent J* 2006;201:37-42.
205. Boggess KA, Edelstein BL. Oral health in women during preconception and pregnancy: implications for birth outcomes and infant oral health. *Matern Child Health J* 2006;10:169-74.
206. William V, Messer LB, Burrow MF. Molar incisor hypomineralization: review and recommendations for clinical management. *Pediatr Dent* 2006;28:224-32.
207. Weinstein P, Harrison R, Benton T. Motivating mothers to prevent caries: confirming the beneficial effect of counseling. *J Am Dent Assoc* 2006;137:789-93.
208. Du MQ, Tai BJ, Jiang H, Lo EC, Fan MW, Bian Z. A two-year randomized clinical trial of chlorhexidine varnish on dental caries in Chinese preschool children. *J Dent Res* 2006;85:557-9.
209. Brazzelli M, McKenzie L, Fielding S, Fraser C, Clarkson J, Kilonzo M, Waugh N. Systematic review of the effectiveness and cost-effectiveness of HealOzone for the treatment of occlusal pit/fissure caries and root caries. *Health Technol Assess*. 2006;10:80.
210. Bankole OO, Aderinokun GA, Odenloye O, Adeyemi AT. Weaning practices among some Nigerian women: implication on oral health. *Odontostomatol Trop* 2006;29:15-21.
211. Chung MH, Kaste LM, Koerber A, Fadavi S, Punwani I. Dental and medical students' knowledge and opinions of infant oral health. *J Dent Educ* 2006;70:511-7.
212. Liu JF. Effects of Nd:YAG laser pulpotomy on human primary molars. The burden of oral ill health for children. *J Endod* 2006;32:404-7.
213. Nunn JH. *Arch Dis Child* 2006;91:251-3.
214. El-Housseiny A, Farsi N. The effectiveness of two antibacterial regimens on salivary mutans streptococci and lactobacilli in children. *J Clin Pediatr Dent* 2005;30:145-51.
215. Gussy MG, Waters EG, Walsh O, Kilpatrick NM. Early childhood caries: current evidence for aetiology and prevention. *J Paediatr Child Health* 2006;42:37-43.
216. Hugoson A, Koch G, Göthberg C, Helkimo AN, Lundin SA, Norderyd O, Sjödin B, Sondell K. Oral health of individuals aged 3-80 years in Jönköping, Sweden during 30 years (1973-2003). II. Review of clinical and radiographic findings. *Swed Dent J* 2005;29:139-55.

217. Weintraub JA, Ramos-Gomez F, Jue B, Shain S, Hoover CI, Featherstone JD, Gansky SA. Fluoride varnish efficacy in preventing early childhood caries. *J Dent Res* 2006;85:172-6.
218. Poulsen S, Laurberg L, Vaeth M, Jensen U, Haubek D. A field trial of resin-based and glass-ionomer fissure sealants: clinical and radiographic assessment of caries. *Community Dent Oral Epidemiol* 2006;34:36-40.
219. Vachirarojpisan T, Shinada K, Kawaguchi Y. The process and outcome of a programme for preventing early childhood caries in Thailand. *Community Dent Health* 2005;22:253-9.
220. Ziegler P, Briefel R, Clusen N, Devaney B. Feeding Infants and Toddlers Study (FITS): development of the FITS survey in comparison to other dietary survey methods. *J Am Diet Assoc* 2006;106:S12-27.
221. Tavener JA, Davies GM, Davies RM, Ellwood RP. The prevalence and severity of fluorosis in children who received toothpaste containing either 440 or 1,450 ppm F from the age of 12 months in deprived and less deprived communities. *Caries Res* 2006;40:66-72.
222. Vann WF Jr, Bouwens TJ, Braithwaite AS, Lee JY. The childhood obesity epidemic: a role for pediatric dentists? *Pediatr Dent* 2005;27:271-6.
223. Pekiner FN, Yücelten D, Ozbayrak S, Sezen EC. Oral-clinical findings and management of epidermolysis bullosa. *J Clin Pediatr Dent* 2005;30:59-65.
224. Pinheiro SL, Simionato MR, Imparato JC, Oda M. Antibacterial activity of glass-ionomer cement containing antibiotics on caries lesion microorganisms. *Am J Dent* 2005;18:261-6.
225. Huth KC, Paschos E, Brand K, Hickel R. Effect of ozone on non-cavitated fissure carious lesions in permanent molars. A controlled prospective clinical study. *Am J Dent* 2005;18:223-8.
226. Balciuniene I, Sabalaite R, Juskiene I. Chemomechanical caries removal for children. *Stomatologija* 2005;7:40-4.
227. Baghdadi ZD. The clinical evaluation of a single-bottle adhesive system with three restorative materials in children: six-month results. *Gen Dent* 2005;53:357-65.
228. Sá Roriz Fonteles C, Zero DT, Moss ME, Fu J. Fluoride concentrations in enamel and dentin of primary teeth after pre- and postnatal fluoride exposure. *Caries Res* 2005;39:505-8.
229. Pascon FM, Kantovitz KR, Caldo-Teixeira AS, Borges AF, Silva TN, Puppini-Rontani RM, Garcia-Godoy F. Clinical evaluation of composite and compomer restorations in primary teeth: 24-month results. *J Dent* 2006;34:381-8.
230. Paschos E, Huth KC, Benz C, Reeka-Bardschmidt A, Hickel R. Efficacy of intraoral topical anesthetics in children. *J Dent* 2006;34:398-404.
231. Plotzitz B, Kneist S, Berger J, Hetzer G. Efficacy of chlorhexidine varnish applications in the prevention of early childhood caries. *Eur J Paediatr Dent* 2005;6:149-54.
232. Wennhall I, Mårtensson EM, Sjunnesson I, Matsson L, Schröder U, Twetman S. Caries-preventive effect of an oral health program for preschool children in a low socio-economic, multicultural area in Sweden: results after one year. *Acta Odontol Scand* 2005;63:163-7.
233. Bergmann J, Leitão J, Kultje C, Bergmann D, Clode MJ. Removing dentine caries in deciduous teeth with Carisolv: a randomised, controlled, prospective study with six-month follow-up, comparing chemomechanical treatment with drilling. *Oral Health Prev Dent* 2005;3:105-11.
234. Tinanoff N. Association of diet with dental caries in preschool children. *Dent Clin North Am* 2005;49:725-37.
235. Kitchens DH. The economics of pit and fissure sealants in preventive dentistry: a review. *J Contemp Dent Pract* 2005;15:6:95-103.
236. Thaweboon S, Thaweboon B, Soo-Ampon S, Soo-Ampon M. Salivary mutans streptococci and lactobacilli after self arresting caries treatment. *Southeast Asian J Trop Med Public Health* 2005;36:765-8.
237. Gisselsson H, Emilson CG, Birkhed D, Björn AL. Approximal caries increment in two cohorts of schoolchildren after discontinuation of a professional flossing program with chlorhexidine gel. *Caries Res* 2005;39:350-6.
238. Wong MC, Lam KF, Lo EC. Bayesian analysis of clustered interval-censored data. *J Dent Res* 2005;84:817-21.
239. Davies GM, Duxbury JT, Boothman NJ, Davies RM, Blinkhorn AS. A staged intervention dental health promotion programme to reduce early childhood caries. *Community Dent Health* 2005;22:118-22.
240. Truin GJ, van't Hof MA. Caries prevention by professional fluoride gel application on enamel and dentinal lesions in low-caries children. *Caries Res* 2005;39:236-40.

241. Mäkinen KK, Isotupa KP, Mäkinen PL, Söderling E, Song KB, Nam SH, Jeong SH. Six-month polyol chewing-gum programme in kindergarten-age children: a feasibility study focusing on mutans streptococci and dental plaque. *Int Dent J* 2005;55:81-8.
242. Chadwick BL, Treasure ET. Primary care research: difficulties recruiting preschool children to clinical trials. *Int J Paediatr Dent* 2005;15:197-204.
243. Türksel Dülgergil C, Satici O, Yildirim I, Yavuz I. Prevention of caries in children by preventive and operative dental care for mothers in rural Anatolia, Turkey. *Acta Odontol Scand* 2004;62:251-7.
244. Thorild I, Lindau B, Twetman S. Salivary mutans streptococci and dental caries in three-year-old children after maternal exposure to chewing gums containing combinations of xylitol, sorbitol, chlorhexidine, and fluoride. *Acta Odontol Scand* 2004;62:245-50.
245. Doufexi A, Mina M, Ioannidou E. Gingival overgrowth in children: epidemiology, pathogenesis, and complications. A literature review. *J Periodontol* 2005;76:3-10.
246. Chan KM, King NM, Kilpatri NM. Can infants catch caries? A review of the current evidence on the infectious nature of dental caries in infants. *N Z Dent J* 2005;101:4-11.
247. Berg JH. Good evidence for incremental preventive benefit of topical fluoride therapies. *Arch Pediatr Adolesc Med* 2005;159:315-6.
248. Jackson RJ, Newman HN, Smart GJ, Stokes E, Hogan JI, Brown C, Seres J. The effects of a supervised toothbrushing programme on the caries increment of primary school children, initially aged 5-6 years. *Caries Res* 2005;39:108-15.
249. Ramalingam L, Messer LB. Early childhood caries: an update. *Singapore Dent J* 2004;26:21-9.
250. Jiang H, Bian Z, Tai BJ, Du MQ, Peng B. The effect of a bi-annual professional application of APF foam on dental caries increment in primary teeth: 24-month clinical trial. *J Dent Res* 2005;84:265-8.
251. Thorild I, Lindau B, Twetman S. Effect of maternal use of chewing gums containing xylitol, chlorhexidine or fluoride on mutans streptococci colonization in the mothers' infant children. *Oral Health Prev Dent* 2003;1:53-7.
252. Twetman S, Stecksén-Blicks C. Effect of xylitol-containing chewing gums on lactic acid production in dental plaque from caries active pre-school children. *Oral Health Prev Dent* 2003;1:195-9.
253. Enwonwu CO, Phillips RS, Ibrahim CD, Danfillo IS. Nutrition and oral health in Africa. *Int Dent J* 2004;54:344-51.
254. Douglass JM, Douglass AB, Silk HJ. A practical guide to infant oral health. *Am Fam Physician*. 2004;70:2113-20.
255. Chadwick BL, Treasure ET, Playle RA. A randomised controlled trial to determine the effectiveness of glass ionomer sealants in pre-school children. *Caries Res* 2005;39:34-40.
256. Graves CE, Berkowitz RJ, Proskin HM, Chase I, Weinstein P, Billings R. Clinical outcomes for early childhood caries: influence of aggressive dental surgery. *J Dent Child (Chic)* 2004;71:114-7.
257. Marx J, Pretorius E. Asthma--a risk factor for dental caries. *SADJ* 2004;59:323, 325-6.
258. Kotsanos N, Dionysopoulos P. Lack of effect of fluoride releasing resin modified glass ionomer restorations on the contacting surface of adjacent primary molars. a clinical prospective study. *Eur J Paediatr Dent* 2004;5:136-42.
259. Turgut MD, Tekçiçek M, Olmez S. Clinical evaluation of a polyacid-modified resin composite under different conditioning methods in primary teeth. *Oper Dent* 2004;29:515-23.
260. Tavener JA, Davies GM, Davies RM, Ellwood RP. The prevalence and severity of fluorosis and other developmental defects of enamel in children who received free fluoride toothpaste containing either 440 or 1450 ppm F from the age of 12 months. *Community Dent Health* 2004;21:217-23.
261. Nainar SM, Mohammed S. Diet counseling during the infant oral health visit. *Pediatr Dent*. 2004;26:459-62.
262. Gaffney KE, Farrar-Simpson MA, Claire D, Davilla G. Prolonged baby bottle feeding: a health risk factor. *Pediatr Nurs* 2004;30:242-5.
263. Weinstein P, Harrison R, Benton T. Motivating parents to prevent caries in their young children: one-year findings. *J Am Dent Assoc* 2004;135:731-8.
264. Attari N, Roberts JF. Restoration of primary teeth affected by Early Childhood Caries. *Eur J Paediatr Dent* 2004;5:92-7.
265. De Grauwe A, Aps JK, Martens LC. Early Childhood Caries (ECC): what's in a name? *Eur J Paediatr Dent* 2004;5:62-70.
266. Ellwood RP, Davies GM, Worthington HV, Blinkhorn AS, Taylor GO, Davies RM. Relationship between area deprivation and the anticaries benefit of an oral health programme providing free fluoride toothpaste to young children. *Community Dent Oral Epidemiol* 2004;32:159-65.

267. Steiner M, Helfenstein U, Menghini G. Effect of 1000 ppm relative to 250 ppm fluoride toothpaste. A meta-analysis. *Am J Dent* 2004;17:85-8.
268. Eight-year study on conventional glass ionomer and amalgam restorations in primary teeth. Qvist V, Laurberg L, Poulsen A, Teglers PT. *Acta Odontol Scand* 2004;62:37-45.
269. Bader JD, Rozier RG, Lohr KN, Frame PS. Physicians' roles in preventing dental caries in preschool children: a summary of the evidence for the U.S. Preventive Services Task Force. *Am J Prev Med* 2004;26:315-25.
270. do Rego MA, Koga-Ito CY, Jorge AO. Effects of oral environment stabilization procedures on counts of *Candida* spp. in children. *Pesqui Odontol Bras* 2003;17:332-6.
271. Casas MJ, Kenny DJ, Johnston DH, Judd PL. Long-term outcomes of primary molar ferric sulfate pulpotomy and root canal therapy. *Pediatr Dent* 2004;26:44-8.
272. Kavvadia K, Karagianni V, Polychronopoulou A, Papagiannouli L. Primary teeth caries removal using the Carisolv chemomechanical method: a clinical trial. *Pediatr Dent* 2004;26:23-8.
273. Amin MS, Harrison RL, Benton TS, Roberts M, Weinstein P. Effect of povidone-iodine on *Streptococcus mutans* in children with extensive dental caries. *Pediatr Dent* 2004;26:5-10.
274. Pine CM, Adair PM, Nicoll AD, Burnside G, Petersen PE, Beighton D, Gillett A, Anderson R, Anwar S, Brailsford S, Broukal Z, Chestnutt IG, Declerck D, Ping FX, Ferro R, Freeman R, Gugushe T, Harris R, Lin B, Lo EC, Maupomé G, Moola MH, Naidoo S, Ramos-Gomez F, Samaranayake LP, Shahid S, Skeie MS, Splieth C, Sutton BK, Soo TC, Whelton H. International comparisons of health inequalities in childhood dental caries. *Community Dent Health* 2004;21:121-30.
275. Pine CM, Adair PM, Burnside G, Nicoll AD, Gillett A, Borges-Yáñez SA, Broukal Z, Brown J, Declerck D, Ping FX, Gugushe T, Hunsrisakhun J, Lo EC, Naidoo S, Nyandindi U, Poulsen VJ, Razanamihaja N, Splieth C, Sutton BK, Soo TC, Whelton H. Barriers to the treatment of childhood caries perceived by dentists working in different countries. *Community Dent Health* 2004;21:112-20.
276. Adair PM, Pine CM, Burnside G, Nicoll AD, Gillett A, Anwar S, Broukal Z, Chestnutt IG, Declerck D, Ping FX, Ferro R, Freeman R, Grant-Mills D, Gugushe T, Hunsrisakhun J, Irigoyen-Camacho M, Lo EC, Moola MH, Naidoo S, Nyandindi U, Poulsen VJ, Ramos-Gomez F, Razanamihaja N, Shahid S, Skeie MS, Skur OP, Splieth C, Soo TC, Whelton H, Young DW. Familial and cultural perceptions and beliefs of oral hygiene and dietary practices among ethnically and socio-economically diverse groups. *Community Dent Health* 2004;21:102-11.
277. Beighton D, Brailsford S, Samaranayake LP, Brown JP, Ping FX, Grant-Mills D, Harris R, Lo EC, Naidoo S, Ramos-Gomez F, Soo TC, Burnside G, Pine CM. A multi-country comparison of caries-associated microflora in demographically diverse children. *Community Dent Health* 2004;21:96-101.
278. Pine CM, Adair PM, Petersen PE, Douglass C, Burnside G, Nicoll AD, Gillett A, Anderson R, Beighton D, Jin-You B, Broukal Z, Brown JP, Chestnutt IG, Declerck D, Devine D, Espelid I, Falcolini G, Ping FX, Freeman R, Gibbons D, Gugushe T, Harris R, Kirkham J, Lo EC, Marsh P, Maupomé G, Naidoo S, Ramos-Gomez F, Sutton BK, Williams S. Developing explanatory models of health inequalities in childhood dental caries. *Community Dent Health* 2004;21:86-95.
279. Harris R, Nicoll AD, Adair PM, Pine CM. Risk factors for dental caries in young children: a systematic review of the literature. *Community Dent Health* 2004;21:71-85.
280. Qvist V, Laurberg L, Poulsen A, Teglers PT. Class II restorations in primary teeth: 7-year study on three resin-modified glass ionomer cements and a compomer. *Eur J Oral Sci* 2004;112:188-96.
281. Nainar SM, Mohammed S. Role of infant feeding practices on the dental health of children. *Clin Pediatr (Phila)* 2004;43:129-33.
282. Petersen PE, Peng B, Tai B, Bian Z, Fan M. Effect of a school-based oral health education programme in Wuhan City, Peoples Republic of China. *Int Dent J* 2004;54:33-41.
283. Rong WS, Bian JY, Wang WJ, Wang JD. Effectiveness of an oral health education and caries prevention program in kindergartens in China. *Community Dent Oral Epidemiol* 2003;31:412-6.
284. Trairatvorakul C, Piwat S. Comparative clinical evaluation of slot versus dovetail Class III composite restorations in primary anterior teeth. *J Clin Pediatr Dent* 2004;28:125-9.
285. Kovari H, Pienihäkkinen K, Alanen P. Use of xylitol chewing gum in daycare centers: a follow-up study in Savonlinna, Finland. *Acta Odontol Scand* 2003;61:367-70.

286. Twetman S, Axelsson S, Dahlgren H, Holm AK, Källestål C, Lagerlöf F, Lingström P, Mejäre I, Nordenram G, Norlund A, Petersson LG, Söder B. Caries-preventive effect of fluoride toothpaste: a systematic review. *Acta Odontol Scand* 2003;61:347-55.
287. Källestål C, Norlund A, Söder B, Nordenram G, Dahlgren H, Petersson LG, Lagerlöf F, Axelsson S, Lingström P, Mejäre I, Holm AK, Twetman S. Economic evaluation of dental caries prevention: a systematic review. *Acta Odontol Scand* 2003;61:341-6.
288. Ruottinen S, Karjalainen S, Pienihäkkinen K, Lagström H, Niinikoski H, Salminen M, Rönnemaa T, Simell O. Sucrose intake since infancy and dental health in 10-year-old children. *Caries Res* 2004;38:142-8.
289. van Rijkom HM, Truin GJ, van 't Hof MA. Caries-inhibiting effect of professional fluoride gel application in low-caries children initially aged 4.5-6.5 years. *Caries Res* 2004;38:115-23.
290. Nourallah AW, Splieth CH. Efficacy of occlusal plaque removal in erupting molars: a comparison of an electric toothbrush and the cross-toothbrushing technique. *Caries Res* 2004;38:91-4.
291. Kaakko T, Skaret E, Getz T, Hujoel P, Grembowski D, Moore CS, Milgrom P. An ABCD program to increase access to dental care for children enrolled in Medicaid in a rural county. *J Public Health Dent* 2002;62:45-50.
292. Davies GM, Worthington HV, Ellwood RP, Blinkhorn AS, Taylor GO, Davies RM, Considine J. An assessment of the cost effectiveness of a postal toothpaste programme to prevent caries among five-year-old children in the North West of England. *Community Dent Health* 2003;20:207-10.
293. Curzon ME, Preston AJ. Risk groups: nursing bottle caries/caries in the elderly. *Caries Res* 2004;38:24-33.
294. Seminario AL, Ivancaková R. Early childhood caries. *Acta Medica (Hradec Kralove)* 2003;46:91-4.
295. Seow WK. Diagnosis and management of unusual dental abscesses in children. *Aust Dent J* 2003;48:156-68.
296. Brook I. Microbiology and management of endodontic infections in children. *J Clin Pediatr Dent* 2003;28:13-7.
297. Blinkhorn AS, Gratrix D, Holloway PJ, Wainwright-Stringer YM, Ward SJ, Worthington HV. A cluster randomised, controlled trial of the value of dental health educators in general dental practice. *Br Dent J* 2003;195:395-400.
298. Lewis CW, Milgrom P. Fluoride. *Pediatr Rev* 2003;24:327-36.
299. Touger-Decker R, van Loveren C. Sugars and dental caries. *Am J Clin Nutr* 2003;78:881S-892S.
300. Zanata RL, Navarro MF, Pereira JC, Franco EB, Lauris JR, Barbosa SH. Effect of caries preventive measures directed to expectant mothers on caries experience in their children. *Braz Dent J* 2003;14:75-81.
301. Krol DM. Dental caries, oral health, and pediatricians. *Curr Probl Pediatr Adolesc Health Care* 2003;33:253-70.
302. Herasse M, Spentchian M, Taillandier A, Keppler-Noreuil K, Florito AN, Bergoffen J, Wallerstein R, Muti C, Simon-Bouy B, Mornet E. Molecular study of three cases of odontohypophosphatasia resulting from heterozygosity for mutations in the tissue non-specific alkaline phosphatase gene. *J Med Genet* 2003;40:605-9.
303. Davies RM, Davies GM, Ellwood RP. Prevention. Part 4: Toothbrushing: what advice should be given to patients? *Br Dent J* 2003;195:135-41.
304. Chussid S. Optimizing infant and toddler oral health. The importance of early intervention. *Dent Today* 2003;22:122-5.
305. Espelid I, Mejäre I, Weerheijm K; EAPD. EAPD guidelines for use of radiographs in children. *Eur J Paediatr Dent* 2003;4:40-8.
306. Vanderas AP, Skamnakis J. Effectiveness of preventive treatment on approximal caries progression in posterior primary and permanent teeth: a review. *Eur J Paediatr Dent* 2003;4:9-15.
307. Poulsen P. Retention of glassionomer sealant in primary teeth in young children. *Eur J Paediatr Dent* 2003;4:96-8.
308. Bian JY, Wang WH, Wang WJ, Rong WS, Lo EC. Effect of fluoridated milk on caries in primary teeth: 21-month results. *Community Dent Oral Epidemiol* 2003;31:241-5.
309. Silegy T, Jacks ST. Pediatric oral conscious sedation. *J Calif Dent Assoc* 2003;31:413-8.
310. Nainar SM, Straffon LH. Targeting of the year one dental visit for United States children. *Int J Paediatr Dent* 2003;13:258-63.
311. Tinanoff N, Palmer CA. Dietary determinants of dental caries and dietary recommendations for preschool children. *Refuat Hapeh Vehashinayim* 2003;20:8-23, 78.
312. Welbury RR, Thomason JM, Fitzgerald JL, Steen IN, Marshall NJ, Foster HE. Increased prevalence of dental caries and poor oral hygiene in juvenile idiopathic arthritis. *Rheumatology (Oxford)* 2003;42:1445-51.
313. Wandera M, Twa-Twa J. Baseline survey of oral health of primary and secondary school pupils in Uganda. *Afr Health Sci* 2003;3:19-22.

314. Foley J, Blackwell A. In vivo cariostatic effect of black copper cement on carious dentine. *Caries Res* 2003;37:254-60.
315. Rivera N, Reyes E, Mazzaoui S, Morón A. Pulpal therapy for primary teeth: formocresol vs electrosurgery: a clinical study. *J Dent Child (Chic)* 2003;70:71-3.
316. Levy SM. An update on fluorides and fluorosis. *J Can Dent Assoc* 2003;69:286-91.
317. González MC, Ruíz JA, Fajardo MC, Gómez AD, Moreno CS, Ochoa MJ, Rojas LM. Comparison of the def index with Nyvad's caries diagnostic criteria in 3- and 4-year-old Colombian children. *Pediatr Dent* 2003;25:132-6.
318. Featherstone JD, Adair SM, Anderson MH, Berkowitz RJ, Bird WF, Crall JJ, Den Besten PK, Donly KJ, Glassman P, Milgrom P, Roth JR, Snow R, Stewart RE. Caries management by risk assessment: consensus statement, April 2002. *J Calif Dent Assoc* 2003;31:257-69.
319. Ammari AB, Bloch-Zupan A, Ashley PF. Systematic review of studies comparing the anti-caries efficacy of children's toothpaste containing 600 ppm of fluoride or less with high fluoride toothpastes of 1,000 ppm or above. *Caries Res* 2003;37:85-92.
320. Bönecker M, Cleaton-Jones P. Trends in dental caries in Latin American and Caribbean 5-6- and 11-13-year-old children: a systematic review. *Community Dent Oral Epidemiol* 2003;31:152-7.
321. DenBesten P, Berkowitz R. Early childhood caries: an overview with reference to our experience in California. *J Calif Dent Assoc* 2003;31:139-43.
322. Berkowitz RJ. Acquisition and transmission of mutans streptococci. *J Calif Dent Assoc* 2003;31:135-8.
323. Featherstone JD. The caries balance: contributing factors and early detection. *J Calif Dent Assoc* 2003;31:129-33.
324. Korhonen M, Käkilehto T, Larmas M. Tooth-by-tooth survival analysis of the first caries attack in different age cohorts and health centers in Finland. *Acta Odontol Scand* 2003;61:1-5.
325. Peressini S. Pacifier use and early childhood caries: an evidence-based study of the literature. *J Can Dent Assoc* 2003;69:16-9.
326. Hübel S, Mejère I. Conventional versus resin-modified glass-ionomer cement for Class II restorations in primary molars. A 3-year clinical study. *Int J Paediatr Dent* 2003;13:2-8.
327. Schriks MC, van Amerongen WE. Atraumatic perspectives of ART: psychological and physiological aspects of treatment with and without rotary instruments. *Community Dent Oral Epidemiol* 2003;31:15-20.
328. Tinanoff N, Kanellis MJ, Vargas CM. Current understanding of the epidemiology mechanisms, and prevention of dental caries in preschool children. *Pediatr Dent* 2002;24:543-51.
329. Li Y, Dasanayake AP, Caufield PW, Elliott RR, Butts JT 3rd. Characterization of maternal mutans streptococci transmission in an African American population. *Dent Clin North Am* 2003;47:87-101.
330. Massara ML, Alves JB, Brandão PR. Atraumatic restorative treatment: clinical, ultrastructural and chemical analysis. *Caries Res* 2002;36:430-6.
331. Scheifele E, Studen-Pavlovich D, Markovic N. Practitioner's guide to fluoride. *Dent Clin North Am* 2002;46:831-46.
332. Waggoner WF. Restoring primary anterior teeth. *Pediatr Dent* 2002;24:511-6.
333. Lee JK. Restoration of primary anterior teeth: review of the literature. *Pediatr Dent* 2002;24:506-10.
334. Seale NS. The use of stainless steel crowns. *Pediatr Dent* 2002;24:501-5.
335. Croll TP, Nicholson JW. Glass ionomer cements in pediatric dentistry: review of the literature. *Pediatr Dent* 2002;24:423-9.
336. Chu CH, Lo EC, Lin HC. Effectiveness of silver diamine fluoride and sodium fluoride varnish in arresting dentin caries in Chinese pre-school children. *J Dent Res* 2002;81:767-70.
337. Edelstein BL. Dental care considerations for young children. *Spec Care Dentist* 2002;22:11S-25S.
338. Duggal MS, Tumba KJ, Sharma NK. Clinical performance of a compomer and amalgam for the interproximal restoration of primary molars: a 24-month evaluation. *Br Dent J* 2002;193:339-42.
339. Davies GM, Worthington HV, Ellwood RP, Bentley EM, Blinkhorn AS, Taylor GO, Davies RM. A randomised controlled trial of the effectiveness of providing free fluoride toothpaste from the age of 12 months on reducing caries in 5-6 year old children. *Community Dent Health* 2002;19:131-6.
340. Cumow MM, Pine CM, Burnside G, Nicholson JA, Chesters RK, Huntington E. A randomised controlled trial of the efficacy of supervised toothbrushing in high-caries-risk children. *Caries Res* 2002;36:294-300.
341. Dasanayake AP, Wiener HW, Li Y, Vermund SH, Caufield PW. Lack of effect of chlorhexidine varnish on *Streptococcus mutans* transmission and caries in mothers and children. *Caries Res* 2002;36:288-93.

342. Autio JT. Effect of xylitol chewing gum on salivary *Streptococcus mutans* in preschool children. *ASDC J Dent Child* 2002;69:81-6, 13.
343. You BJ, Jian WW, Sheng RW, Jun Q, Wa WC, Bartizek RD, Biesbrock AR. Caries prevention in Chinese children with sodium fluoride dentifrice delivered through a kindergarten-based oral health program in China. *J Clin Dent* 2002;13:179-84.
344. Falster CA, Araujo FB, Straffon LH, Nör JE. Indirect pulp treatment: in vivo outcomes of an adhesive resin system vs calcium hydroxide for protection of the dentin-pulp complex. *Pediatr Dent* 2002;24:241-8.
345. Lopez L, Berkowitz R, Spiekerman C, Weinstein P. Topical antimicrobial therapy in the prevention of early childhood caries: a follow-up report. *Pediatr Dent* 2002;24:204-6.
346. Mandel ID. Oral infections: impact on human health, well-being, and health-care costs. *Compend Contin Educ Dent* 2002;23:403-6, 408, 410.
347. Farazi AM, Lupi-Pegurier L, Muller M, Weerheijm KL. Restorative materials used on primary teeth: a comparative study between two European universities: Nice and Amsterdam. *ASDC J Dent Child* 2001;68:339-43, 301.
348. Peretz B. Juvenile dermatomyositis: literature review and report of a case. *Compend Contin Educ Dent* 2001;22:858-62, 864.
349. Moore PA, Cuddy MA, Magera JA, Caputo AC, Chen AH, Wilkinson LA. Oral transmucosal fentanyl pretreatment for outpatient general anesthesia. *Anesth Prog* 2000;47:29-34.
350. Dutta BN, Gauba K, Tewari A, Chawla HS. Silver amalgam versus resin modified GIC class-II restorations in primary molars: twelve month clinical evaluation. *J Indian Soc Pedod Prev Dent* 2001;19:118-22.
351. Lo EC, Chu CH, Lin HC. A community-based caries control program for pre-school children using topical fluorides: 18-month results. *J Dent Res* 2001;80:2071-4.
352. Sheehy EC, Brailsford SR, Kidd EA, Beighton D, Zoitopoulos L. Comparison between visual examination and a laser fluorescence system for in vivo diagnosis of occlusal caries. *Caries Res* 2001;35:421-6.
353. Näse L, Hatakka K, Savilahti E, Saxelin M, Pönkä A, Poussa T, Korpela R, Meurman JH. Effect of long-term consumption of a probiotic bacterium, *Lactobacillus rhamnosus* GG, in milk on dental caries and caries risk in children. *Caries Res* 2001;35:412-20.
354. Duggal MS, van Loveren C. Dental considerations for dietary counselling. *Int Dent J* 2001;51:408-12.
355. Blinkhorn AS, Wainwright-Stringer YM, Holloway PJ. Dental health knowledge and attitudes of regularly attending mothers of high-risk, pre-school children. *Int Dent J* 2001;51:435-8.
356. Slade GD. Epidemiology of dental pain and dental caries among children and adolescents. *Community Dent Health* 2001;18:219-27.
357. Burt BA, Pai S. Does low birthweight increase the risk of caries? A systematic review. *J Dent Educ* 2001;65:1024-7.
358. Shenkin JD, Davis MJ, Corbin SB. The oral health of special needs children: dentistry's challenge to provide care. *ASDC J Dent Child* 2001;68:201-5.
359. Gomez SS, Weber AA, Emilson CG. A prospective study of a caries prevention program in pregnant women and their children five and six years of age. *ASDC J Dent Child* 2001;68:191-5, 152.
360. Wendt LK, Carlsson E, Hallonsten AL, Birkhed D. Early dental caries risk assessment and prevention in pre-school children: evaluation of a new strategy for dental care in a field study. *Acta Odontol Scand* 2001;59:261-6.
361. Autio-Gold JT, Courts F. Assessing the effect of fluoride varnish on early enamel carious lesions in the primary dentition. *J Am Dent Assoc* 2001;132:1247-53.
362. Attin T, Opatowski A, Meyer C, Zingg-Meyer B, Buchalla W, Mönting JS. Three-year follow up assessment of Class II restorations in primary molars with a polyacid-modified composite resin and a hybrid composite. *Am J Dent* 2001;14:148-52.
363. Wilson NH. Conference report. Direct adhesive materials: current perceptions and evidence--future solutions. *J Dent* 2001;29:307-16.
364. Yalman N, Sepet E, Aren G, Mete Z, Külekçi G, Anak S. The effect of bone marrow transplantation on systemic and oral health in Fanconi's aplastic anemia. *J Clin Pediatr Dent* 2001;25:329-32.
365. Casamassimo PS. Maternal oral health. *Dent Clin North Am* 2001;45:469-78.
366. Vanobbergen J, Martens L, Declerk D. Caries prevalence in Belgian children: a review. *Int J Paediatr Dent* 2001;11:164-70.

367. Boneta AE, Neesmith A, Mankodi S, Berkowitz HJ, Sánchez L, Mostler K, Stewart B, Sintes J, De Vizio W, Petrone ME, Volpe AR, Zhang YP, McCool JJ, Bustillo E, Proskin HM. The enhanced anticaries efficacy of a sodium fluoride and dicalcium phosphate dihydrate dentifrice in a dual-chambered tube. A 2-year caries clinical study on children in the United States of America. *Am J Dent* 2001;14:13A-17A.
368. Cleaton-Jones P. Dental caries trends in 5 to 6 year-old and 11 to 13 year-old children in two UNICEF designated regions: Sub-Saharan Africa, and Middle East and North Africa, 1970-2000. *Refuat Hapeh Vehashinayim* 2001;18:11-22, 75.
369. Lin YT, Tsai CL. Comparative anti-caries effects of tablet and liquid fluorides in cleft children. *J Clin Dent* 2000;11:104-6.
370. Strohenger L, Brambilla E. The use of fluoride varnishes in the prevention of dental caries: a short review. *Oral Dis* 2001;7:71-80.
371. Attrill DC, Ashley PF. Occlusal caries detection in primary teeth: a comparison of DIAGNOdent with conventional methods. *Br Dent J* 2001;190:440-3.
372. More rigorous studies needed to advance emerging dental caries diagnostic and management strategies, says NIH consensus panel. National Institutes of Health Consensus Development Conference. Diagnosis and Management of Dental Caries Throughout Life. New release. *Pediatr Dent* 2001;23:123-4.
373. Wight NE. Management of common breastfeeding issues. *Pediatr Clin North Am* 2001;48:321-44.
374. Willershausen B, Watermann L. Longitudinal study to assess the effectivity of electric and manual toothbrushes for children. *Eur J Med Res* 2001;6:39-45.
375. Waterhouse PJ, Nunn JH, Whitworth JM, Soames JV. Primary molar pulp therapy--histological evaluation of failure. *Int J Paediatr Dent* 2000;10:313-21.
376. Gomez SS, Weber AA. Effectiveness of a caries preventive program in pregnant women and new mothers on their offspring. *Int J Paediatr Dent* 2001;11:117-22.
377. Ramos-Gomez FJ, Petru A, Hilton JF, Canchola AJ, Wara D, Greenspan JS. Oral manifestations and dental status in paediatric HIV infection. *Int J Paediatr Dent* 2000;10:3-11.
378. Donaldson M, Kinirons M. Effectiveness of the school dental screening programme in stimulating dental attendance for children in need of treatment in Northern Ireland. *Community Dent Oral Epidemiol* 2001;29:143-9.
379. Chuckpaiwong S, Nakornchai S, Surarit R, Soo-ampon S. Fluoride analysis of human milk in remote areas of Thailand. *Southeast Asian J Trop Med Public Health* 2000;31:583-6.
380. Autio JT, Courts FJ. Acceptance of the xylitol chewing gum regimen by preschool children and teachers in a Head Start program: a pilot study. *Pediatr Dent* 2001;23:71-4.
381. Gross LC, Griffen AL, Casamassimo PS. Compomers as Class II restorations in primary molars. *Pediatr Dent* 2001;23:24-7.
382. Eidelman E, Holan G, Fuks AB. Mineral trioxide aggregate vs. formocresol in pulpotomized primary molars: a preliminary report. *Pediatr Dent* 2001;23:15-8.
383. Williams AC, Bearn D, Mildinhal S, Murphy T, Sell D, Shaw WC, Murray JJ, Sandy JR. Cleft lip and palate care in the United Kingdom--the Clinical Standards Advisory Group (CSAG) Study. Part 2: dentofacial outcomes and patient satisfaction. *Cleft Palate Craniofac J* 2001;38:24-9.
384. Sandy JR, Williams AC, Bearn D, Mildinhal S, Murphy T, Sell D, Murray JJ, Shaw WC. Cleft lip and palate care in the United Kingdom--the Clinical Standards Advisory Group (CSAG) Study. Part 1: background and methodology. *Cleft Palate Craniofac J* 2001;38:20-3.
385. Croll TP, Helpin ML, Donly KJ. Vitremer restorative cement for children: three clinicians' observations in three pediatric dental practices. *ASDC J Dent Child* 2000;67:391-8, 374.
386. Attin T, Opatowski A, Meyer C, Zingg-Meyer B, Mönning JS. Class II restorations with a polyacid-modified composite resin in primary molars placed in a dental practice: results of a two-year clinical evaluation. *Oper Dent* 2000;25:259-64.
387. Valaitis R, Hesch R, Passarelli C, Sheehan D, Sinton J. A systematic review of the relationship between breastfeeding and early childhood caries. *Can J Public Health* 2000;91:411-7.
388. Chu CH. Treatment of early childhood caries: a review and case report. *Gen Dent* 2000;48:142-8.
389. Ayers KM, Colquhoun AN. Leukaemia in children. Part II--Dental care of the leukaemic child, including management of oral side effects of cancer treatment. *N Z Dent J* 2000;96:141-6.

390. Pine CM, McGoldrick PM, Burnside G, Curnow MM, Chesters RK, Nicholson J, Huntington E. An intervention programme to establish regular toothbrushing: understanding parents' beliefs and motivating children. *Int Dent J* 2000;312-23.
391. Aaltonen AS, Suhonen JT, Tenovuoto J, Inkilä-Saari I. Efficacy of a slow-release device containing fluoride, xylitol and sorbitol in preventing infant caries. *Acta Odontol Scand* 2000;58:285-92.
392. Uhari M, Tapiainen T, Kontiokari T. Xylitol in preventing acute otitis media. *Vaccine* 2000;19:144-7.
393. Isokangas P, Söderling E, Pienihäkkinen K, Alanen P. Occurrence of dental decay in children after maternal consumption of xylitol chewing gum, a follow-up from 0 to 5 years of age. *J Dent Res* 2000;79:1885-9.
394. Martof A. Consultation with the specialist: dental care. *Pediatr Rev* 2001;22:13-5.
395. Edelstein BL. Access to dental care for Head Start enrollees. *J Public Health Dent* 2000;60:221-9.
396. Kanellis MJ. Caries risk assessment and prevention: strategies for Head Start, Early Head Start, and WIC. *J Public Health Dent* 2000;60:210-7.
397. Tinanoff N, Palmer CA. Dietary determinants of dental caries and dietary recommendations for preschool children. *J Public Health Dent* 2000;60:197-206.
398. Fomon SJ, Ekstrand J, Ziegler EE. Fluoride intake and prevalence of dental fluorosis: trends in fluoride intake with special attention to infants. *J Public Health Dent* 2000;60:131-9.
399. Edelstein BL. Public and clinical policy considerations in maximizing children's oral health. *Pediatr Clin North Am* 2000;47:1177-89.
400. Nowak AJ, Warren JJ. Infant oral health and oral habits. *Pediatr Clin North Am* 2000;47:1043-66.
401. Schafer TE, Adair SM. Prevention of dental disease. The role of the pediatrician. *Pediatr Clin North Am* 2000;47:1021-42.
402. Caufield PW, Griffen AL. Dental caries. An infectious and transmissible disease. *Pediatr Clin North Am* 2000;47:1001-19.
403. Welbury RR, Shaw AJ, Murray JJ, Gordon PH, McCabe JF. Clinical evaluation of paired compomer and glass ionomer restorations in primary molars: final results after 42 months. *Br Dent J* 2000;189:93-7.
404. Mascarenhas AK. Risk factors for dental fluorosis: a review of the recent literature. *Pediatr Dent* 2000;22:269-77.
405. Clarkson JJ, McLoughlin J. Role of fluoride in oral health promotion. *Int Dent J* 2000;50:119-28.
406. Bucher HU, Baumgartner R, Bucher N, Seiler M, Fauchère JC. Artificial sweetener reduces nociceptive reaction in term newborn infants. *Early Hum Dev* 2000;59:51-60.
407. Nelson M. Childhood nutrition and poverty. *Proc Nutr Soc* 2000;59:307-15.
408. Twetman S, García-Godoy F, Goepferd SJ. Infant oral health. *Dent Clin North Am* 2000;44:487-505.
409. Featherstone JD. The science and practice of caries prevention. *J Am Dent Assoc* 2000;131:887-99.
410. Milgrom P, Reisine S. Oral health in the United States: the post-fluoride generation. *Annu Rev Public Health* 2000;21:403-36.
411. van Lunsen DM, de Soet JJ, Weerheijm KL, Groen HJ, Veerkamp JS. Effects of dental treatment and single application of a 40% chlorhexidine varnish on mutans Streptococci in young children under intravenous anaesthesia. *Caries Res* 2000;34:268-74.

PAPER II

Early Childhood Caries (ECC) and background factors. A case control cross-sectional study.

Congiu Giovanna^a, Lugliè PierFranca^a, Cagetti MariaGrazia^b, Sale Silvana^a, Spano Giovanni^a, Campus Guglielmo^a.

^aDental Institute, University of Sassari, ^bWHO Collaborating Centre of Milan for Epidemiology and Community Dentistry

Corresponding author

Dr Guglielmo Campus

Dental Institute, University of Sassari

Viale san Pietro 43/C I-07100 Sassari, Italy

Tel.+39-079-228540

Fax. +39-079-228541

Email gcampus@uniss.it

Summary

Aim: to evaluate the influence of several determinants, like as socio-economical level (SES), behavioural factors (dietary and oral hygiene) on the presence of ECC in pre-school children in the town of Sassari.

Design: a cross-sectional case control study was designed. The protocol consisted in a dental examination and a standardized questionnaire. The study group was of 544 subjects: 260 girls (47.8%) and 284 boys (52.2%). The sample was then categorized into two age groups: 185 (34.0%) children aged 48-60 months and 359 (66.0%) children aged 18-47 months.

Results: caries experience was 16.0% (28.6% in the older group and 9.5% in the younger one). Caries presence was linked to: a) parents' educational level in both parents in the old groups ($p<0.01$); b) mother's educational level ($p<0.01$) in the young group; c) use of bottle feeding ($p=0.02$) in the older group; d) use of sweetened dummy at night ($p=0.01$) in the young group; e) brushing habits after each meal ($p=0.04$) in the younger group; f) dental check-ups in the last six months ($p=0.01$) in the young group; g) eruption age of the first teeth ($p=0.04$) in the young group; h) number of siblings in both age groups ($p<0.01$). In the conditional logistic regression procedure a dummy variable of the parents' educational level was created. An educational level of parents shows a protective effect towards ECC prevalence ($OR=0.44$ 95%CI 0.34 – 0.57). Also the presence in the family of one or more siblings was statically associated to ECC ($OR=1.91$ 95%CI 1.30 – 2.81).

Conclusion: our results highlight the role played by socio-economic status and behavioural habits in the development of ECC.

During the past decades a severe decrease in caries experience in children in countries of western world (Szoke and Petersen, 2000; Campus *et al.*, 2001; Vanobbergen, 2004; Campus *et al.*, 2007; Campus *et al.*, 2009) was observed. In industrialised countries, the prevalence, severity and patterns of caries are thought to be related to determinants which include social and educational background, dietary and oral hygiene practices (Roeters *et al.*, 1995; Steckslen-Blicks and Borssen, 1999). However, the pattern of caries disease reflects distinct risk profiles related to living conditions, lifestyles and environmental factors and the implementation of preventive oral health schemes (Petersen, 2005). Findings of relationships between socioeconomic status (SES) and health outcomes are quite ubiquitous across in the health literature. People with lower educational attainment and lower income usually have lower life expectancy; consequently, children from families with a low income are more likely to have a low birth weight and to suffer from childhood illnesses (Armfield, 2007). Breastfeeding is recommended by paediatricians to be continued for at least the first year of life and beyond for as long as mutually desired by mother and child (Gartner *et al.*, 2005). Prolonged and unrestricted breastfeeding, however, has been reported to be a potential risk factor for caries (Hallett *et al.*, 2006; Iida *et al.*, 2007). Various systems of classification have been used to define Early Childhood Caries (ECC) (Drury *et al.*, 1999; Psoter *et al.*, 2003). Two commonly accepted classifications for ECC include simple ECC and severe ECC, as defined by the American Academy of Paediatric Dentistry (AAPD, 2008) the first one Simple ECC any deciduous presence of 1 or more decayed, missing, or filled tooth surfaces in any primary tooth in a child under 6 years of age is the most used one.

Caries disease in preschool children has a complex multifactorial aetiology: it has also been associated with improper feeding practice (Ismail, 2003; Hallett *et al.*, 2006;

Campus *et al.*, 2007; Kramer *et al.*, 2007) and low socio-economic background (Ismail, 2003; Muirhead and Marcenes, 2004; Hallett *et al.*, 2006). Nowadays, increasing attention is being paid to differences in population subgroups in characterizing the distribution and correlation of dental disease (Seppä *et al.*, 2000; Muirhead and Marcenes, 2004; Bankel *et al.*, 2006). A link between preterm birth or low-birth weight and caries was proposed (Seow, 1997; Burt and Pai, 2001; Eastman, 2003; Shulman, 2005; Saraiva *et al.*, 2007), but no conclusive evidence has been found.

The present study evaluates the influence of several determinants, like as socio-economical level (SES), behavioural factors (dietary and oral hygiene) on the presence of ECC in pre-school children in the town of Sassari. The null hypothesis was that no associations are present between ECC presence and socio-behavioural factors. To validate this hypothesis a cross-sectional case control study was designed and performed.

Material and methods

Design of study and sample size selection

The study was conducted from January 2010 to November 2010. Data on the total number of residents in Sassari from pre-school groups were derived from the National Statistical Institute (Italian National office of Statistics, ISTAT <http://www.demo.istat>). The total number of children aged 18-60 months was 5354 (2714 boys and 2640 girls). Sassari is the largest district of northern Sardinia with 127.893 inhabitants. The fluoride concentration in local tap water is low (0.03 mg/L) (Campus *et al.*, 2009). The study designed was approved by the ethical committee of the University of Sassari (n°184/2009).

Caries prevalence data on Sardinian preschool children was used (Campus *et al.*, 2009) to calculate sample size. The theoretical sample size with a Confidence Interval set at 5 was 529 children, the number was increased by 5% against the possible number of non-responders. Children were recruited at the kindergartens using systematic cluster sampling, where each class was identified as a cluster and compiled into a list. The first cluster was randomly chosen, while the others were selected at the systematic interval of three classes. Altogether 722 children were recruited in the study (Figure 1).

The protocol consisted in a dental examination and a standardized questionnaire. Parents or guardians were issued with an information leaflet, explaining the aim of the study and requesting their child's participation, after which they were asked for signed consent. Only children with questionnaire filled out and parents signed consent were enrolled in the study (621 subjects). The clinical examination was performed during the schooldays with the teacher/s present. Sixty-one (9.8%) children were absent on the school day of the clinical examination and another 16 (2.6%) refused to participate. The study group was of 544 subjects: 260 girls (47.8%) and 284 boys (52.2%). The sample was then categorized into two age groups: 185 (34.0%) children aged 48-60 months and 359 (66.0%) children aged 18-47 months.

Clinical examination

Clinical assessments were carried out by one examiner (GC) under standardized conditions using a drying tooth device, a plain mirror and a WHO-CPI probe under optimal artificial lighting. Dental caries experience (dmfs) was recorded. A carious lesion was diagnosed when a cavity at dentinal level was noted (Pitts *et al.*, 1995).

The examiner received training and intra- and inter-examiner reliability was assessed before the beginning of the survey. Fifty subjects were examined and re-examined after

72 hours. Inter-examiner reliability was evaluated through analysis of variance for fixed effect compared to a benchmark (Castiglia *et al.*, 2007), while intra-examiner reproducibility was assessed as percent agreement and Cohen's Kappa statistics. A good examiner reliability compared to benchmark was found without significant differences ($p = 0.15$) and with a low value of mean squares for error (0.51). As regard, the percent agreement, intra-examiner reproducibility, was high (Cohen's Kappa 0.86). The standardized questionnaire consisted of closed questions. It was filled out by parents or guardians at home and brought to school by the child. It contained questions related to medical status, consumption of sweets and soft drinks (mean intake frequency/day), frequency of toothbrushing and use of fluoride supplements apart from fluoride toothpaste.

Data analysis

Data were coded and imputed into a database (Microsoft® Excel 2010) and checked to verify accuracy of data. Statistical analysis was performed using Stata® 10.0 for Mac (<http://www.stata.com>). Initially, clinical condition parameters and potential risk indicators were analyzed univariately to describe the variables and distributions. Descriptive statistical analysis was performed. To avoid the attenuating effect of unequal variability among groups on the value of t , a square root transformation was performed when the response variable was a count (Fleiss, 1986). The association between ECC and background factors was tested using χ^2 test. Multivariate analysis was performed at two levels, first a logistic regression forward stepwise procedure using as dependent variable caries experience positively $dmfs > 0$ was run in the two age groups separately. The conditional logistic regression procedure was used to the

analysis of a sample of matched 1:2 case-control. Unless stated otherwise, the criterion for statistical significance was set at $\alpha=0.05$.

Results

Overall caries experience was 16.0% (28.6% in the older group and 9.5% in the younger one). Mean dmfs was 0.9 ± 0.2 $_{95\%CI}$ (0.6 – 1.3) in all subjects while in the older group was 1.9 ± 0.5 $_{95\%CI}$ (1.0 – 2.8) and was 0.4 ± 0.1 $_{95\%CI}$ (0.2 – 0.6). The need of dental care ($d > 0$) was predominant in all age groups. The distribution of demographic and social variables by caries prevalence is displayed in Table 1. The parents' educational level was significantly associated to caries prevalence in both parents in the old groups ($p < 0.01$), in the young group caries prevalence was associated to the mother's educational level ($p < 0.01$). Regarding dietary habits the use of bottle feeding was associated to caries prevalence ($p = 0.02$) in the older group; in the young group the use of sweetened dummy at night was statically associated to caries ($p = 0.01$). In the younger group, brushing habits after each meal ($p = 0.04$), dental check-ups in the last six months ($p = 0.01$), the eruption age of the first teeth ($p = 0.04$) were statistically linked to caries prevalence. The number of siblings was statistical associated in both age groups ($p < 0.01$). Two logistic regression models were constructed and run for each age group separately. A high educational level of the father was protective to the presence of caries lesion in older sample, while the educational level of the mother was statistically significant in the younger group (Table 2). The presence in the family of more than one sibling was positively associated to caries ($OR = 1.63$ $_{95\%CI}$ 1.06 – 2.49). In the conditional logistic regression procedure a dummy variable of the parents' educational level was created (Table 3). An educational level of parents shows a protective effect towards ECC prevalence ($OR = 0.44$ $_{95\%CI}$ 0.34 – 0.57). Also the

presence in the family of one or more siblings was statically associated to ECC (OR=1.91 95%CI 1.30 – 2.81).

Discussion

Caries experience in Sardinian kindergarten children was similar than that reported in other surveys (Sundby and Petersen, 2003), moreover an impressive decrease was observed respect to those previously reported (Campus *et al.*, 2007). As reported in literature, the socio-economical status of the family and behavioural habits has a substantial impact on the development of ECC (Armfield, 2007). Low socio-economic status compromises the ability of individuals to care for their health leading to a reduced resistance to oral and other diseases.

Uncorrected nursing habits (bottle-feeding, use of pacifier at night, etc.) are the most frequently reported causes of ECC (Ismail, 2003), but the disease may also occur in children who are breast-fed. In the United States, for example, the majority of babies are fed using a nursing bottle, yet the majority of them do not develop ECC (Iida *et al.*, 2007).

In Italy, primary dental health service is based on private health care providers; thus, oral care is mainly financed by direct payment or, to a lesser extent, through public or private insurance schemes. The low level of caries experience described in this paper supports the hypothesis that public dental services play a minor role in the caries experience decrease observed in all developed countries.

Several numbers of limitations should be also considered when interpreting these findings. For example the study was cross-sectional and, consequently, no information is available regarding to lesion progression; the methods used for caries diagnosis

(visual inspection with probe and mirror) may be considered a limitation, but these methods are considered to have excellent reliability (Castiglia *et al.*, 2007).

In conclusion, our results highlight the role played by socio-economic status and behavioural habits in the development of ECC. We hope that the results of our study help in the promotion of improved oral health amongst infants and small children in Sassari.

References

- American Academy of Paediatric Dentistry (2008). Definition of Early Childhood Caries (ECC). http://www.aapd.org/media/policies_guidelines/d_ecc.pdf
- Armfield JM. Socioeconomic inequalities in child oral health: a comparison of discrete and composite area-based measures. *J Public Health Dent* 2007;67:119-25.
- Bankel M, Eriksson UC, Robertson A, Köhler B. Caries and associated factors in a group of Swedish children 2- 3 years of age. *Swed Dent J* 2006;30:137-46.
- Burt BA, Pai S. Does low birthweight increase the risk of caries? A systematic review. *J Dent Educ* 2001;65:1024-7.
- Campus G, Cagetti MG, Sacco G, Benedetti G, Strohmer L, Lingström P. Caries risk profiles in Sardinian schoolchildren using Cariogram. *Acta Odontol Scand* 2009;67:146-52.
- Campus G, Lumbau A, Lai S, Solinas G, Castiglia P. Socio-economic and behavioural factors related to caries in twelve-year-old Sardinian children. *Caries Res* 2001;35:427-34.
- Campus G, Solinas G, Sanna A, Maida C, Castiglia P. Determinants of ECC in Sardinian preschool children. *Community Dent Health* 2007;24:253-6.
- Campus G, Solinas G, Strohmer L, Cagetti MG, Sanna A, Minelli L, Majori S, Montagna MT, Reali D, Castiglia P; Collaborating Study Group. National pathfinder survey on children's oral health in Italy: pattern and severity of caries disease in 4-year-olds. *Caries Res* 2009;43:155-62.
- Castiglia P, Campus G, Solinas G, Maida C, Strohmer L. Children's oral health in Italy: training and clinical calibration of examiners for the National Pathfinder about caries disease. *Oral Health Prev Dent* 2007;5:255-61.

Drury TF, Horowitz AM, Ismail AI, Maertens MP, Rozier RG, Selwitz RH. Diagnosing and reporting early childhood caries for research purposes. A report of a workshop sponsored by the National Institute of Dental and Craniofacial Research, the Health Resources and Services Administration, and the Health Care Financing Administration. *J Public Health Dent* 1999;59:192-7.

Eastman DL. Dental outcomes of preterm infants. *Newborn Infant Nurs Rev* 2003;3:93-98.

Fleiss JL. Confidence intervals vs significance tests: quantitative interpretation. *Am J Public Health*. 1986 May;76(5):587-8.

Gartner SL. Breastfeeding. *Am J Nurs* 2005;105:15.

Hallett KB, O'Rourke PK. Caries experience in preschool children referred for specialist dental care in hospital. *Aust Dent J* 2006;51:124-9.

<http://www.stata.com>

Iida H, Auinger P, Billings RJ, Weitzman M. Association between infant breastfeeding and early childhood caries in the United States. *Pediatrics* 2007;120:944-52.

Ismail AI. Determinants of health in children and the problem of early childhood caries. *Pediatr Dent* 2003;25:328-33.

Italian National office of Statistics. ISTAT <http://www.demo.istat>

Kramer MS, Matush L, Vanilovich I, Platt RW, Bogdanovich N, Sevkovskaya Z, Dzikovich I, Shishko G, Collet JP, Martin RM, Davey Smith G, Gillman MW, Chalmers B, Hodnett E, Shapiro S; PROBIT Study Group. Effects of prolonged and exclusive breastfeeding on child height, weight, adiposity, and blood pressure at age 6.5 y: evidence from a large randomized trial. *Am J Clin Nutr* 2007;86:1717-21.

Muirhead V, Marcenes W. An ecological study of caries experience, school performance and material deprivation in 5-year-old state primary school children. *Community Dent Oral Epidemiol* 2004;32:265-70.

Petersen PE, Bourgeois D, Bratthall D, Ogawa H. Oral health information systems--towards measuring progress in oral health promotion and disease prevention. *Bull World Health Organ* 2005;83:686-93.

Pitts NB, Longbottom C. Preventive Care Advised (PCA)/Operative Care Advised (OCA) - categorising caries by the management option. *Community Dent Oral Epidemiol* 1995;23:55-9.

Psoter WJ, Zhang H, Pendrys DG, Morse DE, Mayne ST. Classification of dental caries patterns in the primary dentition: a multidimensional scaling analysis. *Community Dent Oral Epidemiol* 2003;31:231-8.

Roeters J, Burgersdijk R, Truin GJ, van 't Hof M. Dental caries and its determinants in 2-to-5-year-old children. *ASDC J Dent Child* 1995;62:401-8.

Saraiva MC, Chiga S, Bettiol H, Silva AA, Barbieri MA. Is low birthweight associated with dental caries in permanent dentition? *Paediatr Perinat Epidemiol* 2007;21:49-56.

Seow WK. Effects of preterm birth on oral growth and development. *Aust Dent J* 1997;42:85-91.

Seppä L, Kärkkäinen S, Hausen H. Caries trends 1992-1998 in two low-fluoride Finnish towns formerly with and without fluoridation. *Caries Res* 2000;34:462-8.

Shulman JD. Is there an association between low birth weight and caries in the primary dentition? *Caries Res* 2005;39:161-7.

Stecksén-Blicks C, Borssén E. Dental caries, sugar-eating habits and toothbrushing in groups of 4-year-old children 1967-1997 in the city of Umeå, Sweden. *Caries Res* 1999;33:409-14.

Szöke J, Petersen PE. Evidence for dental caries decline among children in an East European country (Hungary). *Community Dent Oral Epidemiol* 2000;28:155-60.

Vanobbergen J, Declerck D, Mwalili S, Martens L. The effectiveness of a 6-year oral health education programme for primary schoolchildren. *Community Dent Oral Epidemiol* 2004; 32:173-82.

Table 1. Distribution of Demographic and Social Variables by caries prevalence (simple-ECC).

	dmfs=0 (n %)	dmfs>0 (n %)	Person chi-sqaure p-value
Sample n=544	457 (84.0)	87 (16.0)	
males n=284 (52.2)	237 (83.4)	47 (16.6)	
females n=260 (47.8)	220 (84.6)	40 (15.4)	
48-60 months			
males n=100	76 (76.0)	24 (24.0)	
females n=85	56 (65.8)	29 (34.2)	$\chi^2=1.66$ p=0.54
18-47 months			
males n=184	161 (87.5)	23 (12.5)	
females n=175	164 (93.7)	11 (6.3)	$\chi^2=4.04$ p=0.04
<i>Mother's educational level</i>			
48-60 months (no responders =5)			
• Primary schools	36 (20.0)	28 (15.6)	
• Secondary schools	62 (34.4)	20 (11.1)	
• University degree	30 (16.7)	4 (2.2)	$\chi^2=12.54$ p<0.01
18-47 months (no responders =6)			
• Primary schools	64 (17.8)	14 (3.9)	
• Secondary schools	138 (38.4)	14 (3.9)	
• University degree	123 (34.3)	6 (1.7)	$\chi^2=10.04$ p<0.01
<i>Father's educational level</i>			
48-60 months			
• Primary schools	52 (29.0)	36 (20.1)	
• Secondary schools	44 (24.6)	13 (7.3)	
• University degree	31 (17.3)	3 (1.7)	$\chi^2=13.83$ p<0.01
18-47 months			
• Primary schools	95 (27.0)	15 (4.3)	
• Secondary schools	129 (36.8)	11 (3.1)	
• University degree	94 (26.8)	7 (2.0)	$\chi^2=3.43$ p=0.180
<i>Mother's occupational status</i>			
48-60 months (no responders =3)			
• Unemployed	42 (23.0)	26 (14.3)	
• Employed	88 (48.4)	26 (14.3)	$\chi^2=4.96$ p=0.03
18-47 months (no responders =0)			
• Unemployed	62 (17.4)	14 (4.0)	
• Employed	261 (73.3)	19 (5.3)	$\chi^2=9.62$ p<0.01
<i>Father's occupational status</i>			
48-60 months (no responders =5)			
• Unemployed	6 (3.3)	7 (3.9)	
• Employed	78 (43.4)	29 (16.1)	
• Self working	45 (25.0)	15 (8.3)	$\chi^2=4.57$ p=0.10
18-47 months (no responders =7)			
• Unemployed	18 (5.1)	7 (2.0)	
• Employed	191 (54.3)	18 (5.1)	
• Self working	109 (31.0)	9 (2.5)	$\chi^2=10.45$ p<0.01
<i>Sweetened dummy at night</i>			
48-60 months (no responders =4)			
• Yes	26 (14.4)	14 (7.7)	
• No	104 (57.5)	37 (20.4)	$\chi^2=1.18$ p=0.27
18-47 months (no responders =1)			
• Yes	56 (15.7)	12 (3.3)	
• No	268 (74.9)	22 (6.1)	$\chi^2=6.48$ p=0.01

<i>Bottle feeding</i>			
48-60 months (no responders =8)			
• Yes	111 (62.7)	34 (19.2)	$\chi^2=5.46$ p=0.02
• No	18 (10.2)	14 (7.9)	
18-47 months (no responders =10)			
• Yes	260 (74.5)	26 (7.5)	$\chi^2=0.01$ p=0.91
• No	57 (16.3)	6 (1.7)	
<hr/>			
<i>Meals a day</i>			
48-60 months (no responders =4)			
• 3	27 (14.9)	10 (5.5)	$\chi^2=5.32$ p=0.07
• 4	69 (38.1)	19 (10.5)	
• >4	34 (18.8)	22 (12.2)	
18-47 months (no responders =5)			
• 3	35 (9.9)	5 (1.4)	$\chi^2=3.43$ p=0.18
• 4	199 (56.2)	15 (4.2)	
• >4	87 (24.6)	13 (3.7)	
<hr/>			
<i>Dental check-ups</i>			
48-60 months (no responders =4)			
• Yes	32 (17.7)	15 (8.3)	$\chi^2=0.31$ p=0.57
• No	97 (53.6)	37 (20.4)	
18-47 months (no responders =2)			
• Yes	20 (5.6)	6 (1.7)	$\chi^2=5.97$ p=0.01
• No	303 (84.9)	28 (7.8)	
<hr/>			
<i>Eruption of the first teeth</i>			
48-60 months (no responders =13)			
• 6 months	47 (27.3)	16 (9.3)	$\chi^2=0.11$ p=0.94
• 9 months	46 (26.7)	17 (9.9)	
• >9 months	33 (19.2)	13 (7.6)	
18-47 months (no responders =12)			
• 6 months	105 (30.3)	18 (5.2)	$\chi^2=6.07$ p=0.04
• 9 months	123 (35.4)	10 (2.9)	
• >9 months	86 (24.8)	5 (1.4)	
<hr/>			
<i>Brush teeth after meal</i>			
48-60 months (no responders =2)			
• Yes	48 (26.3)	20 (10.9)	$\chi^2=0.01$ p=0.91
• No	82 (44.8)	33 (18.0)	
18-47 months (no responders =4)			
• Yes	51 (14.4)	10 (2.8)	$\chi^2=3.95$ p=0.04
• No	270 (76.0)	24 (6.8)	
<hr/>			
<i>Number of siblings</i>			
48-60 months (no responders =6)			
• 0	48 (26.2)	13 (7.1)	$\chi^2=17.49$ p<0.01
• 1	41 (22.4)	6 (3.3)	
• >1	41 (22.4)	34 (18.6)	
18-47 months (no responders =2)			
• 0	108 (30.2)	5 (1.4)	$\chi^2=9.53$ p<0.01
• 1	67 (18.8)	4 (1.1)	
• >1	148 (41.5)	25 (7.0)	

Table 2. Logistic estimates of the model (forward stepwise logistic regression) for caries experience, by age groups.

48-60 months

Number of observations = 178

Log Likelihood = 89.43, $\chi^2_4 = 30.54$ $P < 0.01$

Variable	<i>P</i>	OR	CI _{95%}
Mother's educational level	0.06	0.55	0.30-1.03
Father's educational level	0.02	0.48	0.26-0.90
Meals a day	0.05	1.66	1.00-2.77
Number of siblings	0.02	1.63	1.06-2.49

18-47 months

Number of observations = 349

Log Likelihood = -167.32, $\chi^2_4 = 33.70$ $P < 0.01$

Variable	<i>P</i>	OR	CI _{95%}
Mother's educational level	<0.01	0.52	0.36-0.77
Father's occupational status	0.08	0.65	0.40-1.05
Sweetened dummy at night	0.07	0.56	0.30-1.06
Number of siblings	0.10	1.23	0.96-1.59


Table 3. Odds ratio (OR) and 95% confidence intervals for ECC.

Number of observations = 239

Log Likelihood = -111.69, $\chi^2_4 = 69.12$ $P < 0.01$

	Odds Ratio	95% Confidence Interval
Parents' educational level	0.44	0.34 – 0.57
Meals a day	1.62	0.99 – 2.64
Number of siblings	1.91	1.30 – 2.81

Figure 1. Flow chart of the study sample.



PAPER III

Fluoride concentration in breast milk after the use of a fluoridated food supplement. A randomized placebo-controlled study.

Guglielmo Campus^{ab}, Giovanna Congiu^a, Silvana Sale^a, Maria Grazia Cagetti^b, Gavino Sanna^c, Peter Lingstrom^{bd}

^aDental Institute, University of Sassari, ^bWHO Collaborating Centre of Milan for Epidemiology and Community Dentistry, ^cChemical Department , University of Sassari, Italy, ^dDepartment of Cariology, Institute of Odontology at Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden.

Corresponding author
Dr Guglielmo Campus
Dental Institute, University of Sassari
Viale san Pietro 43/C I-07100 Sassari, Italy
Tel.+39-079-228540
Fax. +39-079-228541
Email gcampus@uniss.it

Summary

Background A way to administer fluoride to infants is to provide fluoride to the breastfeeding mother, but no sufficient scientific evidence is available.

Aim: The hypothesis behind this report was that the use of a fluoride food supplement in mothers during breastfeeding might increase the fluoride concentration in breast milk.

Design: A double-blind randomized placebo-controlled study was designed. 124 women were divided in two groups: a fluoride group, using a non-sucrose food supplement containing fluoride and a control group, using a non-sucrose food supplement without fluoride. The investigation had a randomized placebo-controlled study design with an experimental period of six weeks.

Results: at t_0 , fluoride concentration values in both groups were quite comparable and no statistically significant differences were observed ($p=0.52$ respectively). The dropout rate was quite high (40.3%) mainly caused by the early interruption of breastfeeding. The comparison of the fluoride concentrations at the three different endpoints (t_0 - t_1 - t_2) between the two groups did not show statistical differences ($p=0.12$).

Conclusions No statistically significant differences were found between fluoride group and control group. Overall, the fluoride concentrations in the two groups did not significantly changed during the experimental period; only a statistically significant increase in F concentration was noted in the fluoride group.

Despite great efforts and achievements in oral health promotion were obtained, dental caries still remains a major health problem in childhood (Petersen, 2009). A striking reduction in dental caries in industrialized countries was registered, although the proportion with severe disease has remained at 10-15%, and the prevalence is increasing in developing countries. In Italy as in the majority of industrialized countries, recent data display an uneven distribution, with the highest burden of the disease among the underprivileged groups (Campus *et al.*, 2007; 2009; Wennhall *et al.*, 2008); this distribution calls for new strategies in caries prevention.

Milk and milk products contain nutrients that have potential anticaries properties: calcium, phosphate, casein, and lipids (Levine, 2001; Aimutis, 2004). A low prevalence of caries among schoolchildren who drink milk daily was recorded; the sample did not use fluoride regularly and had poor oral hygiene (Petti *et al.*, 1997). However, results of epidemiological studies on the association between dairy products intake and dental caries among young children are scant and inconsistent. A large debate was carried on the efficacy of fluoride during pregnancy. Some studies demonstrated fluoride ions capacity to pass the placental barrier (Toyama *et al.*, 2001; Gupta *et al.*, 1993); however still now scientific evidence on the possibility of fluoride ion to be incorporated into the child's teeth during pregnancy period is not available. A way to administer fluoride to infants is to provide fluoride to the lactating mother. The amount of fluoride in breast milk has been studied for many years (Sener *et al.*, 2007; Chuckpaiwong *et al.*, 2000; Latifah *et al.*, 1989). The fluoride concentration in breast milk was evaluated in women living in areas with low and high fluoride concentration in water (Spak *et al.*, 1983), but no statistically significant difference was found. The hypothesis behind this report was that the use of a fluoride food supplement in lactating mothers increases the fluoride concentration in breast milk.

Materials and methods

Study design and sample selection

A double-blind randomized placebo-controlled study was designed and approved by the ethical committee of the University of Sassari (n°184/2009). The study was conducted in Sassari, Sardinia, Italy, between October and December 2009. All the participants were resident in Sassari area, where a low natural fluoride content (≈ 0.3 ppm) in tap water is present. In 2008, the number of delivery was 1041. Women were recruited at the Gynaecologic Clinic of the University of Sassari before the delivery. Using systematic cluster sampling, women were compiled into a list; the first cluster was randomly chosen, while the others were selected at the systematic interval of five women. Subjects with a history of systemic antibiotic, topical fluoride or chlorhexidine treatment within 30 days before the baseline visit were excluded. Inclusion criterium was the possibility to use a breast pump (electric or manual).

Altogether 324 women were recruited. An information leaflet, explaining the aim of the study was delivered and a signed consent was requested. Only women with signed consent were called for examination (286 subjects). The flow chart of the study design is shown in Figure 1. In order to obtain interpretable results, the number of subjects to be included in each group was calculated. Two groups of women were formed: a fluoride group, using a non-sucrose food supplement containing fluoride and a control group, using a non-sucrose food supplement without fluoride. Considering significant a 25% difference between groups in fluoride concentration in breast milk, with values of 40% and 15% for the fluoride and the control groups respectively, and a 95% probability to obtain a significant difference between groups at 5% level, the number of subjects needed per group was 26.5. The acceptance rate was 55.8% (n = 160, mean

age 32.5 ± 4.6). Using a computer program (Excel[®] 2003 for Mac OsX), the randomization was carried out on an individual basis by GC. 36 women refused to participate, thus the final study sample was formed by 124 subjects (63 women in the fluoride group and 61 in the control one). At t_1 interim evaluation, 27 women were excluded due to the early interruption of breastfeeding, (9 in the fluoride group and 18 in the control group), at t_2 , 21 more women were excluded (10 received a systemic antibiotics therapy, 3 did not return the empty blisters and 8 due to the early interruption of breastfeeding). Thus, only 74 women concluded the experimental period: 40 in the fluoride group and 34 in the control group.

Treatment and sample collection

The two supplementations, powder form, were identical in weight (2.17g), form, colour and packing, but only one contained fluoride (1.5 ppm for each dose). They were produced and supplied by Milte[®] Italia spa (Italy), and coded as “green” or “red”. The code was sealed by an independent monitor and not broken until the statistical analysis was finalized. The women were instructed to mix the powder into 100/150 ml of tap water and to drink the solution once a day in the morning after breakfast. Subjects were asked to make no changes in dietary and oral hygiene habits. All subjects received a fluoridated toothpaste containing 1450 $\mu\text{g/g}$ NaF (Mentadent P, Unilever Italia, Milano) to be used during the experimental period. They were asked to avoid any other oral hygiene adjuvant and any commercial fluoride products during the study. Moreover women were instructed to drink bottled water with a low fluoride content (<0.04 ppm). In order to evaluate the success of administration of the food supplement, women were given food supplement necessary for a single week at a time and they were asked to return the empty blisters. Samples of milk were collected at baseline (t_0), after three weeks of food supplement use (t_1) and finally after six weeks of use (t_2).

Milk samples were gathered during 3 minutes by women into a polyethylene tube, after cleaning the breast with cotton wool and distilled water. An aliquot of this milk (5.0µl) was collected in a plastic tip and stored in a sterile plastic microbeaker.

Fluoride analysis

Prior to analysis, an appropriate volume of Total Ionic Strength Adjustment Buffer (TISAB, usually 0.5 ml) was added to all samples, in order to adjust the pH 5.0 and their ionic strength. The fluoride concentrations of the samples were measured after adding 5 ml of distilled water. The chemical analyses were done in a blind manner; fluoride was analyzed using a ion-specific electrode (model 9609, Orion Research) coupled to a potentiometer (model 710A, Orion Research). All measurements were made in triplicate and expressed as mean \pm standard error (SE). A complete speciation protocol was carried out to determine the different fluoride fractions: a- Inorganic Free Fluoride (IFF), b- Inorganic Bonded Fluoride (IBF), c- Casein Organic fluoride (OCF), d- Globulinic Organic Fluoride (OGF) and finally e- Total fluoride concentration (TF).

Statistical Analysis

To normalize the data, fluoride concentrations were logarithmically transformed prior to analysis. Mean, standard deviation and standard error were calculated for each product and phase. The data were multiplied for 1100 (the dilution factor) to obtain the concentration of fluoride. Data were analyzed for statistically significant differences using repeated one-way measures analysis of variance (ANOVA). Anova was calculated after each milk collection and at the end of the research as the sum of fluoride concentrations at all the time intervals, as a single measure of substantivity.

Results

No adverse effect was referred by from women in both groups. A total of 74 women completed the experimental period (Figure 1) and data on fluoride concentration in breast milk are referred to 40 subjects from fluoride group and 34 from control group. The dropout rate was quite high (40.3%), mainly caused by the early interruption of breastfeeding.

The different fluoride fractions recorded at t_2 in the two groups are reported in Table 1. The fluoride concentration in the different fraction were quite similar in the two groups; only in Casein Organic Fluoride (OCF) fraction a slight difference was observed (141 $\mu\text{g/l}$ in the fluoride group and 132 $\mu\text{g/l}$ in the control group) ($p=0.06$).

At t_0 , fluoride concentrations in both groups were quite comparable and no statistically significant differences were observed ($p = 0.52$ respectively).

At t_2 , women in the fluoride group showed a statistically significant increase in fluoride concentration compared to baseline (from $472 \pm 118 \mu\text{g/l}$ to $510 \pm 123 \mu\text{g/l}$) ($p=0.03$). In the control group, a slight not statistically significant increase in fluoride content was observed (from $480 \pm 148 \mu\text{g/l}$ to $486 \pm 150 \mu\text{g/l}$). The comparison of the fluoride concentrations at the three different endpoints (t_0 - t_1 - t_2) between the two groups did not show statistical differences ($p=0.12$) (Table 2).

Discussion

The aim of the present study was to evaluate whether a fluoride food supplement administered to lactating women can increase the fluoride content in breast milk. A randomized clinical trial was designed and carried out. No statistically significant differences were found between fluoride group and control group. Overall, the fluoride

concentrations in the two groups did not significantly changed during the experimental

period. This result can be explained with the existence of a plasma-milk barrier against fluoride, in order to protect the infant from excessive intake of fluoride (Ekstrand *et al.*, 1981). Nevertheless, in the present paper, a statistically significant increase in the halogen concentration was noted in the fluoride group, after six weeks of fluoridated food supplement use.

Breast milk possesses unique nutritional, biochemical, anti-infective and anti-allergic properties. From theoretical point of view, breast milk might represent an excellent way to administer fluoride to infants. A wide range of fluoride concentration in breast milk is reported, *i.e.* WHO reports a range from < 0.002 to 0.1 ppm (World Health Organization). In the present paper the fluoride concentrations measured were higher than those reported in literature (Sener *et al.*, 2007). A strength point of the paper is the inclusion criteria like the recent stipulated absence of recent use of fluoride supplements as a preventive measure. However it is necessary to underline that the large number of drop-out might have affected the study validity and it is possible to speculate that a larger sample could lead to different results.

References

Aimutis WR. Bioactive properties of milk proteins with particular focus on anticariogenesis. *Journal of Nutrition* 2004;134:989S–95S.

Campus G, Solinas G, Sanna A, Maida C, Castiglia P. Determinants of ECC in Sardinian preschool children. *Community Dent Health* 2007;24:253-56.

Campus G, Solinas G, Strohmenger L, Cagetti MG, Senna A, Minelli L, Majori S, Montagna MT, Reali D, Castiglia P, Collaborating Study Group. National pathfinder survey on children's oral health in Italy: pattern and severity of caries disease in 4-year-olds. *Caries Res* 2009;43:155-62.

Chuckpaiwong S, Nakornchai S, Surarit R, Soo-ampon S. Fluoride analysis of human milk in remote areas of Thailand. *Southeast Asian J Trop Med Public Health* 2000;31:583-6.

Gupta S, Seth AK, Gupta A, Gavane AG. Transplacental passage of fluorides. *J Pediatr* 1993;123:139-41.

J Ekstrand, L O Boreus, and P de Chateau. No evidence of transfer of fluoride from plasma to breast milk. *Br Med J (Clin Res Ed)* 1981;283(6294):761–762.

Latifah R, Razak IA. Fluoride levels in mother's milk. *J Pedod* 1989;13:149-54.

Levine RS. Milk, flavoured milk products and caries. *British Dental Journal* 2001;191:20.

Petersen PE. Global policy for improvement of oral health in the 21st century--implications to oral health research of World Health Assembly 2007, World Health Organization. *Community Dent Oral Epidemiol* 2009;37:1-8.

Petti S, Simonetti R, Simonetti D'Arca A. The effect of milk and sucrose consumption on caries in 6-to-11-year-old Italian schoolchildren. *European Journal of Epidemiology* 1997;13:659–64.

Dr.ssa Giovanna Congiu EARLY CHILDHOOD CARIES (ECC) IS STILL A PROBLEM? AN EPIDEMIOLOGICAL AND AETIOLOGICAL APPROACH.
Università degli Studi di Sassari

Sener Y, Tosun G, Kahvecioglu F, Gokalp A, Hasan Koc H. Fluoride Levels of Human Plasma and Breast Milk. *Eur J Dent* 2007;1:21-24.

Spak CJ, Hardell LI, De Chateau P. Fluoride in human milk. *Acta Paediatr Scand* 1983;72:699-701.

Toyama Y, Nakagaki H, Kato S, Huang S, Mizutani Y, Kojima S, Toyama A, Ohno N, Tsuchiya T, Kirkham J, Robinson C. Fluoride concentrations at and near the neonatal line in human deciduous tooth enamel obtained from a naturally fluoridated and a non-fluoridated area. *Arch Oral Biol* 2001;46:147-53.

Wennhall I, Matsson L, Schroder U, Twetman S. Outcome of an oral health outreach programme for preschool children in a low socio-economic multicultural area. *Int J Paediatr Dent* 2008;18:84–90.

World Health Organization. *Environmental Health Criteria*. Geneva: 2002. p. 227.

Figure 1. Flow chart of the study sample.




Table 1. Different Fluoride fractions concentrations in breast milk at t₂.

	Inorganic Free Fluoride (IFF) <i>mean range</i> (µg/l)	Inorganic Bonded Fluoride (IBF) <i>mean range</i> (µg/l)	Casein Organic fluoride (OCF) <i>mean</i> <i>range</i> (µg/l)	Globulinic Organic Fluoride (OGF) <i>mean range</i> (µg/l)	Total fluoride concentration (TF) <i>mean range</i> (µg/l)
Fluoride group	73 (14 ÷ 77)	125 (53 ÷ 185)	141 (75 ÷ 208)	171 (70 ÷ 235)	510 (390 ÷ 620)
Control group	68 (40 ÷ 85)	118 (68 ÷ 140)	132 (88 ÷ 163)	168 (142 ÷ 256)	486 (413 ÷ 604)
<i>Anova one way</i>	NS	NS	0.06	NS	NS

Table 2. Fluoride concentrations in breast milk measured in the two groups at the three different endpoints.

	<i>Fluoride group</i> <i>mean± Standard</i> <i>Deviation (µg/l)</i>	<i>Control group</i> <i>mean± Standard</i> <i>Deviation (µg/l)</i>	<i>Anova one</i> <i>way</i>
t ₀ baseline	472±118	480±148	NS
t ₁ three weeks of use	480±126	472±142	NS
t ₂ six weeks of use	510±123	486±150	
<i>Anova one way</i>	0.03	NS	